

An analytic theory of extremal hypergraph problems*

Vladimir Nikiforov[†]

May 14, 2013

Abstract

The starting point of this paper is the following problem: If $\mathcal{P}$ is a hereditary property of r -uniform graphs, find the limit

$$\pi(\mathcal{P}) = \lim_{n \rightarrow \infty} \binom{n}{r}^{-1} \max\{e(G) : G \in \mathcal{P} \text{ and } v(G) = n\}.$$

It is shown that this problem is just a particular case of a general analytic problem about a parameter $\lambda^{(\alpha)}(G)$ defined for every r -graph G and every real $\alpha \geq 1$ as

$$\lambda^{(\alpha)}(G) = \max_{|x_1|^\alpha + |x_2|^\alpha + \dots + |x_n|^\alpha = 1} r! \sum_{\{i_1, i_2, \dots, i_r\} \in E(G)} x_{i_1} x_{i_2} \dots x_{i_r},$$

Note that $\lambda^{(1)}(G)$ is a well-studied parameter, however, the truly exceptional value is $\lambda(G) = \lambda^{(r)}(G)$, known as the largest eigenvalue of G .

Two of the main results of the paper are: for all $\alpha \geq 1$ the limit

$$\lambda^{(\alpha)}(\mathcal{P}) = \lim_{n \rightarrow \infty} n^{r/\alpha - r} \max\{\lambda^{(\alpha)}(G) : G \in \mathcal{P} \text{ and } v(G) = n\}$$

exists, and if $\alpha > 1$, then

$$\lambda^{(\alpha)}(\mathcal{P}) = \pi(\mathcal{P}).$$

It is shown also that if $\lambda^{(1)}(\mathcal{P}) = \pi(\mathcal{P})$, then $\mathcal{P}$ has remarkable features regarding extremal problems. Many known concrete results are generalized and further research is outlined.

1 Introduction and main results

In this paper we build an analytic theory of hypergraph extremal problems of the type:

Given an r -uniform graph G of order n , belonging to some hereditary property $\mathcal{P}$, how many edges can G have?

One of the key results proved in this paper states that this combinatorial problem is asymptotically equivalent to an extremal analytic problem:

If G is as above, how large the largest eigenvalue of G can be?

***AMS MSC:** 05C65; 05C35. **Keywords:** uniform hypergraphs; extremal problems; hereditary property; largest eigenvalue; graph Lagrangians.

[†]Department of Mathematical Sciences, University of Memphis, Memphis TN 38152, USA; email: vnikifrv@memphis.edu

To clarify this point, let us lay down the definition of the largest eigenvalue $\lambda(G)$ that is adopted here:

Suppose that the vertices of G are the integers $1, 2, \dots, n$ and let $x_1, x_2, \dots, x_n$ be real numbers. Set

$$\lambda(G) = \max_{|x_1|^r + \dots + |x_n|^r = 1} r! \sum \{x_{i_1} x_{i_2} \cdots x_{i_r} : \{i_1, i_2, \dots, i_r\} \text{ is an edge of } G\}. \quad (1)$$

The value $\lambda(G)$ turns out to be at the meeting point of two major lines of research - one is on maxima of homogenous polynomial forms on graphs, and the other is on spectra of hypermatrices. We start by highlighting a few milestones along these two lines, some of which are all but forgotten.

1.1 Background

The study of polynomial forms on graphs and their maxima over unit spheres in the l^1 norm has been pioneered by Motzkin and Straus in [22], and later generalized by Khadziivanov [17] and by Sós and Straus [38], see also [29] for some historical remarks. For hypergraphs the same topic has been studied first by Brown and Simonovits [1], and later by Sidorenko [37], with some very definite results; other similar early contributions are by Frankl and Rödl [7] and Frankl and Füredi [8]. While for 2-graphs this method has been enlightening, for hypergraphs the obtained results were less encompassing, due to the fact that this research remained bound to unit spheres in the l^1 norm, as was the original result of Motzkin and Straus.

On the other hand, for even positive r , the study of critical points of polynomial forms of degree r over finite dimensional unit spheres in the l^r norm has been suggested by Lusternik and Schnirelman already in 1930, see [20], p. 38, or its French translation [21]. This topic has been developed further by Krasnoselskii [18], Elsholtz, Tzitzianadze, and others, but the focal point of these later contributions has shifted to infinite dimensional spaces. Nevertheless, the underlying idea of Lusternik and Schnirelman, forgotten for decades, nowadays became mainstream, under the name of “variational eigenvalues” of hypermatrices. Indeed, in the same spirit, recently Lim [19] proposed a variational approach to spectra of hypermatrices of both even and odd dimensions. Independently, an algebraic approach to the same goal was proposed by Qi in [32]. For further developments on spectra of hypermatrices see [2],[3],[9],[33],[34],[40],[41],[42].

1.2 Overview

One point that needs clarification is why the maximum in (1) is taken over the unit sphere in the l^r norm. First, most of the definitions of eigenvalues adopted in the above cited papers on spectra of hypermatrices reduce precisely to (1) for the largest eigenvalue of hypergraphs. Second, as shown by Friedland, Gauber and Han [9], the use of the l^r norm is indeed necessary, if we want to preserve some essential features of the Perron-Frobenius theory for r -dimensional matrices.

While the largest eigenvalue $\lambda(G)$ is exceptional, in this paper we study a more general parameter $\lambda^{(\alpha)}(G)$, defined exactly as $\lambda(G)$ in (1), but with maximum taken over the unit sphere in the l^α norm, where $\alpha \geq 1$ is a real number. For $\alpha > 1$ the parameter $\lambda^{(\alpha)}(G)$ has been introduced and used by Keevash, Lenz and Mubayi in [16], but they provided just scanty groundwork on $\lambda^{(\alpha)}(G)$. Thus, one of the goals of this paper is to set a more solid base for a systematic study of $\lambda^{(\alpha)}(G)$ and of $\lambda(G)$.

Before continuing let us stress that $\lambda^{(\alpha)}(G)$ is defined as a conditional maximum; thus, its usability in extremal problems is rooted in its very nature. Indeed, for 2-graphs it has been shown that many classical results can be enhanced and recast for the largest eigenvalue $\lambda(G)$ with an astonishing preservation of detail; for a survey of these results see [27], and for some new developments also [31]. Unfortunately, for hypergraphs the present situation is not so advanced: there are just a few isolated spectral extremal results, mainly due to Cooper and Dutle [4] and to Keevash, Lenz and Mubayi [16].

Moreover, $\lambda^{(\alpha)}(G)$ is just one of the many critical values that can be defined in a setting similar to (1), and at least some of them are applicable to extremal graph problems. For example, these possibilities have been explored for the smallest and the second largest eigenvalue of 2-graphs. In this paper we focus exclusively on $\lambda^{(\alpha)}(G)$, with a brief discussion of other similar parameters in the concluding remarks.

As already said, we shall show that edge extremal problems are asymptotically equivalent to extremal problems for the largest eigenvalue. In this way, known extremal edge results readily imply spectral bounds for hypergraphs. However, the opposite implication seems more significant, as it paves the road for an extensive use of differential calculus in hypergraphs. For example, finding the maximum possible eigenvalue of a graph with a forbidden subgraph gives asymptotically the maximum possible number of its edges, yet solving the former problem might be easier, using known analytical techniques.

We also illustrate the use of various proof tools in solving concrete problems, in particular, in problems for flat and multiplicative properties, to be defined later. These tools include the Hypergraph Removal Lemma, classical inequalities, Lagrange multipliers and other methods from real analysis. The concrete results that we obtain shed new light on several older results of Sidorenko [37], and on some new ones by Keevash, Lenz and Mubayi [16].

The paper ends up with a summary discussion and open problems.

1.3 The basics

Recall that an **r -uniform hypergraph** (**r -graph**) consists of a set of vertices $V(G)$ and a set $E(G)$ of r -subsets of V , called edges. We set $v(G) = |V|$ and $e(G) = |E(G)|$. When $V(G)$ is not defined explicitly, it is assumed that $V(G) = [n] = \{1, 2, \dots, n\}$.

Given an r -graph G and a vector $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$, the **polyform** of G is the function $P_G(\mathbf{x})$ defined as

$$P_G(\mathbf{x}) = r! \sum_{\{i_1, i_2, \dots, i_r\} \in E} x_{i_1} x_{i_2} \cdots x_{i_r}.$$

Note that $P_G(\mathbf{x})$ is a homogenous polynomial of degree r and is linear in each variable x_i . Clearly, the definition (1) is equivalent to

$$\lambda(G) = \max_{\|\mathbf{x}\|_r=1} P_G(\mathbf{x}).$$

The largest eigenvalue $\lambda(G)$ has turned out to be a versatile parameter, with close relations to many properties of G ; see [30] for some results. As already said, the choice of the l^r norm in the definition of $\lambda(G)$ makes it exceptional, but it will be useful to consider a more general parameter $\lambda^{(\alpha)}(G)$, defined for every real number $\alpha \geq 1$ as

$$\lambda^{(\alpha)}(G) = \max_{\|\mathbf{x}\|_\alpha=1} P_G(\mathbf{x}) = \max_{|x_1|^\alpha + \dots + |x_n|^\alpha = 1} r! \sum_{\{i_1, i_2, \dots, i_r\} \in E(G)} x_{i_1} x_{i_2} \cdots x_{i_r}.$$

Note first that $\lambda^{(r)}(G) = \lambda(G)$, and second, that $\lambda^{(1)}(G)$ is another much studied parameter, known as the Lagrangian¹ of G . So $\lambda^{(\alpha)}(G)$ can link $\lambda(G)$ to a large body of previous work on extremal hypergraph problems.

The purpose of the following propositions is twofold: first, to give the reader some insight in the meaning and use of $\lambda^{(\alpha)}(G)$; second, these general results, together with the results in Section 2, set the

¹Let us note that this use of the name *Lagrangian* is at odds with the tradition. Indeed, names as *Laplacian*, *Hessian*, *Gramian*, *Grassmanian*, etc., usually denote a structured object like matrix, operator, or manifold, and not just a single number.

background for more thorough future study of $\lambda^{(\alpha)}(G)$. On more than one occasion we shall see the special role of the case $\alpha = r$.

First, taking the n -vector $\mathbf{x} = (n^{-1/\alpha}, \dots, n^{-1/\alpha})$, we immediately get

$$\lambda^{(\alpha)}(G) \geq P_G(\mathbf{x}) = r!e(G)/n^{r/\alpha}.$$

On the other hand, for $\alpha > 1$, Keevash, Lenz and Mubayi have proved that $\lambda^{(\alpha)}(G) \leq (r!e(G))^{1-1/\alpha}$ in [16], Lemma 5. We shall improve this result in Theorem 21 and Corollary 22 below, which also allow for some additional fine-tuning. Here is the summary of these bounds.

Proposition 1 *If $\alpha \geq 1$ and G is an r -graph of order n , then*

$$r!e(G)/n^{r/\alpha} \leq \lambda^{(\alpha)}(G) \leq (r!e(G))^{1-1/\alpha}. \quad (2)$$

If G contains at least one edge, then $\lambda^{(\alpha)}(G) < (r!e(G))^{1-1/\alpha}$.

Inequalities (2) show that $\lambda^{(\alpha)}(G)$ tends to $r!e(G)$ when $\alpha \rightarrow \infty$. Noting that

$$\lambda^{(\alpha)}(G) = \max_{|y_1|+\dots+|y_n|=1} r! \sum_{\{i_1, i_2, \dots, i_r\} \in E(G)} |y_{i_1}|^{1/\alpha} |y_{i_2}|^{1/\alpha} \dots |y_{i_r}|^{1/\alpha}.$$

it becomes also clear that $\lambda^{(\alpha)}(G)$ is increasing and continuous in α .

Proposition 2 *If $\alpha \geq 1$ and G is an r -graph, then $\lambda^{(\alpha)}(G)$ is increasing and continuous in α , and*

$$\lim_{\alpha \rightarrow \infty} \lambda^{(\alpha)}(G) = r!e(G).$$

A cornerstone bound on $\lambda(G)$ for a 2-graph G , with maximum degree Δ , is the inequality $\lambda(G) \leq \Delta$. For r -graphs this generalizes to $\lambda(G) \leq (r-1)!\Delta$, but if $1 \leq \alpha < r$, there is no analogous bound for $\lambda^{(\alpha)}(G)$, which would be tight. Here is what we can say presently on these relations.

Proposition 3 *Let G be an r -graph of order n , with maximum degree Δ .*

(i) If $\alpha \geq r$, then

$$\lambda^{(\alpha)}(G) \leq \frac{(r-1)!\Delta}{n^{r/\alpha-1}}, \quad (3)$$

with equality holding if and only if G is regular;

(ii) If $1 \leq \alpha < r$, there exist r -graphs G for which (3) fails;

(iii) If $1 \leq \alpha < r$, then $\lambda^{(\alpha)}(G) < (r-1)!\Delta$.

Another cornerstone result about $\lambda(G)$ of a 2-graph G of order n is: $\lambda(G) = 2e(G)/n$ if and only if G is regular. We saw in Proposition 1 that the inequality $\lambda(G) \geq 2e(G)/n$ generalizes seamlessly for $\lambda^{(\alpha)}(G)$ of any r -graph G and any $\alpha \geq 1$, but as shown below the condition for equality becomes quite intricate, even for $r = 2$.

Proposition 4 *If $\alpha \geq 1$ and $\lambda^{(\alpha)}(G) = r!e(G)/n^{r/\alpha}$, then G is regular. If $\alpha \geq r$ and G is regular, then $\lambda^{(\alpha)}(G) = r!e(G)/n^{r/\alpha}$. However, if $1 \leq \alpha < r$, then there exist regular graphs G such that $\lambda^{(\alpha)}(G) > r!e(G)/n^{r/\alpha}$.*

Proposition 2 states that $\lambda^{(\alpha)}(G)$ increases in α . Here are two useful technical statements which give some information how fast $\lambda^{(\alpha)}(G)$ can increase indeed.

Proposition 5 *If $\alpha \geq 1$ and G is an r -graph, then the function*

$$h_G(\alpha) = \lambda^{(\alpha)}(G) n^{r/\alpha}$$

is nonincreasing in α . If G is non-regular, then $h_G(\alpha)$ is decreasing in α .

Proposition 6 *If $\alpha \geq 1$ and G is an r -graph, then the function*

$$f_G(\alpha) = \left(\frac{\lambda^{(\alpha)}(G)}{r!e(G)} \right)^\alpha$$

is nonincreasing in α .

1.4 Graph properties and asymptotics of extremal problems

In this paper extremal graph problems are discussed in the general setting of **properties** of r -graphs, which are just families of r -graphs closed under isomorphisms. Given a property $\mathcal{P}$, we shall write $\mathcal{P}_n$ for the set of all graphs in $\mathcal{P}$ of order n . A property is called **monotone** if it is closed under taking subgraphs, and **hereditary**, if it is closed under taking induced subgraphs. For example, given a set of r -graphs $\mathcal{F}$, the family of all r -graphs that do not contain any $F \in \mathcal{F}$ as a subgraph is a monotone property, denoted by $\text{Mon}(\mathcal{F})$. Likewise, the family of all r -graphs that do not contain any $F \in \mathcal{F}$ as an induced subgraph is a hereditary property, denoted as $\text{Her}(\mathcal{F})$. When $\mathcal{F}$ consists of a single graph F , we shall write $\text{Mon}(F)$ and $\text{Her}(F)$ instead of $\text{Mon}(\{F\})$ and $\text{Her}(\{F\})$.

The extremal problems studied below stem from the following one: *Given a hereditary property $\mathcal{P}$ of r -graphs, find*

$$ex(\mathcal{P}, n) = \max_{G \in \mathcal{P}_n} e(G). \quad (4)$$

If $r = 2$ and $\mathcal{P}$ is a monotone property, sharp asymptotics of $ex(\mathcal{P}, n)$ is known, but general hereditary properties seem to have been shrugged off, although a simple and appealing asymptotic solution also exists, see [31] for details. For $r \geq 3$ the problem has turned out to be generally very hard and has been solved only for very few properties $\mathcal{P}$; see [15] for an up-to-date discussion. An easier asymptotic version of the same problem arises from the following fact, established by Katona, Nemetz and Simonovits [13].

Proposition 7 *If $\mathcal{P}$ is a hereditary property of r -graphs, then the sequence*

$$\left\{ ex(\mathcal{P}, n) \binom{n}{r}^{-1} \right\}_{n=1}^{\infty}$$

is nonincreasing and so the limit

$$\pi(\mathcal{P}) = \lim_{n \rightarrow \infty} ex(\mathcal{P}, n) \binom{n}{r}^{-1}$$

always exists.

Thus, if we find $\pi(\mathcal{P})$, we can obtain $ex(\mathcal{P}, n)$ asymptotically, but even $\pi(\mathcal{P})$ is hard to find for most properties $\mathcal{P}$, in particular, $\pi(\text{Mon}(F))$ is not known for many simple graphs F .

As it turns out, the parameters $\lambda^{(\alpha)}(G)$, and in particular $\lambda(G)$, can be efficient tools for the study of $\pi(\mathcal{P})$. Indeed, given a hereditary property $\mathcal{P}$ of r -graphs, set in analogy to (4)

$$\lambda^{(\alpha)}(\mathcal{P}, n) = \max_{G \in \mathcal{P}_n} \lambda^{(\alpha)}(G).$$

Now choosing $G \in \mathcal{P}_n$ with maximum number of edges, Proposition 1 implies that

$$\lambda^{(\alpha)}(G) \geq r! ex(\mathcal{P}, n) / n^{r/\alpha},$$

and so

$$\lambda^{(\alpha)}(\mathcal{P}, n) \geq r! ex(\mathcal{P}, n) / n^{r/\alpha}. \quad (5)$$

Let us begin with a theorem about $\lambda^{(\alpha)}(G)$, which is similar to Proposition 7.

Theorem 8 *Let $\alpha \geq 1$. If $\mathcal{P}$ is a hereditary property of r -graphs, then the limit*

$$\lambda^{(\alpha)}(\mathcal{P}) = \lim_{n \rightarrow \infty} \lambda^{(\alpha)}(\mathcal{P}, n) n^{r/\alpha - r} \quad (6)$$

exists. If $\alpha = 1$, then $\lambda^{(1)}(\mathcal{P}, n)$ is nondecreasing, and so

$$\lambda^{(1)}(\mathcal{P}, n) \leq \lambda^{(1)}(\mathcal{P}). \quad (7)$$

If $\alpha > 1$, then $\lambda^{(\alpha)}(\mathcal{P})$ satisfies

$$\lambda^{(\alpha)}(\mathcal{P}) \leq \frac{\lambda^{(\alpha)}(\mathcal{P}, n) n^{r/\alpha - 1}}{(n-1)(n-2) \dots (n-r+1)}. \quad (8)$$

Interestingly, Theorem 8 is as important as Proposition 7, and its proof is not too hard either, yet it seems to have been missed even in the much studied case $\alpha = 1$.

Here is an immediate consequence of Theorem 8. From (5) we see that

$$\frac{\lambda^{(\alpha)}(\mathcal{P}, n) n^{(r/\alpha) - 1}}{(n-1)(n-2) \dots (n-r+1)} \geq \frac{ex(\mathcal{P}, n)}{\binom{n}{r}},$$

and letting $n \rightarrow \infty$, we find also that

$$\lambda^{(\alpha)}(\mathcal{P}) \geq \pi(\mathcal{P}). \quad (9)$$

We shall show that, almost always, equality holds in this inequality.

An important property of $\lambda^{(\alpha)}(\mathcal{P})$ is that it is nonincreasing in α . Note the difference with Proposition 2, which states that $\lambda^{(\alpha)}(G)$ is increasing in α for every fixed graph G .

Theorem 9 *If $\mathcal{P}$ is a hereditary property of r -graphs, then $\lambda^{(\alpha)}(\mathcal{P})$ is nonincreasing in $\alpha \geq 1$.*

We deduce Theorem 9 from the following subtler relation, which itself is obtained from Proposition 6.

Proposition 10 *If $\mathcal{P}$ is a hereditary property of r -graphs, and $1 \leq \alpha \leq \beta$, then*

$$\left(\frac{\lambda^{(\beta)}(\mathcal{P})}{\pi(\mathcal{P})} \right)^\beta \leq \left(\frac{\lambda^{(\alpha)}(\mathcal{P})}{\pi(\mathcal{P})} \right)^\alpha. \quad (10)$$

Before concluding this subsection, we shall give an immediate application of Theorem 8, but since it refers to blow-up of an r -graph, let us first we define this concept:

*Given an r -graph H of order h and positive integers $k_1, \dots, k_n$, write $H(k_1, \dots, k_h)$ for the graph obtained by replacing each vertex $v \in V(H)$ with a set U_v of size x_v and each edge $\{v_1, \dots, v_r\} \in E(H)$ with a complete r -partite r -graph with vertex classes $U_{v_1}, \dots, U_{v_r}$. The graph $H(k_1, \dots, k_h)$ is called a **blow-up** of H .*

It is well known (see, e.g., [15], Theorem 2.2) that if H is an r -graph of order h and $H(k_1, \dots, k_h)$ is a fixed blow-up of H , then

$$\pi(\text{Mon}(H)) = \pi(\text{Mon}(H(k_1, \dots, k_h))). \quad (11)$$

It turns out that a similar result holds for $\lambda^{(\alpha)}$:

Theorem 11 *If $\alpha > 1$, H is an r -graph of order h and $H(k_1, \dots, k_h)$ is a fixed blow-up of H , then*

$$\lambda^{(\alpha)}(\text{Mon}(H)) = \lambda^{(\alpha)}(\text{Mon}(H(k_1, \dots, k_h))).$$

Our proof of Theorem 11 is not long, but it is based on the Hypergraph Removal Lemma and other fundamental results about r -graphs. Note also that there are simple examples showing that the theorem does not hold for $\alpha = 1$.

1.5 The equivalence of $\lambda^{(\alpha)}(\mathcal{P})$ and $\pi(\mathcal{P})$

Since $\pi(\mathcal{P})$ and $\lambda^{(\alpha)}(\mathcal{P})$ are defined alike, one anticipates some close relation between them to hold. For example, for any r -graph G of order n , inequality (2) implies that

$$\lambda(G) \geq r!e(G)/n. \quad (12)$$

From (9) we get also that if $\mathcal{P}$ is a hereditary property, then

$$\lambda(\mathcal{P}) \geq \pi(\mathcal{P}). \quad (13)$$

Now, if we know that $\lambda(\mathcal{P}) = \pi(\mathcal{P})$, then for every $\alpha > r$, inequality (9) and Theorem 9 imply that

$$\pi(\mathcal{P}) \leq \lambda^{(\alpha)}(\mathcal{P}) \leq \lambda(\mathcal{P}) = \pi(\mathcal{P}),$$

and so, $\lambda^{(\alpha)}(\mathcal{P}) = \pi(\mathcal{P})$ as well.

It turns out that equality always holds in (13), as stated in the following theorem, which is the central result of the paper:

Theorem 12 *If $\mathcal{P}$ is a hereditary property of r -graphs, then for every $\alpha > 1$,*

$$\lambda^{(\alpha)}(\mathcal{P}) = \pi(\mathcal{P}). \quad (14)$$

It seems that a result of this scope is not available in the literature, even for 2-graphs, so some remarks are due here. First, using (14), every result about $\pi(\mathcal{P})$ of a hereditary property $\mathcal{P}$ gives a result about $\lambda(\mathcal{P})$ as well, so we readily obtain a number of results for the largest eigenvalue of uniform graphs. But equality (14) is more significant, as the left and right hand sides of (12) could be quite different. Moreover, finding $\pi(\mathcal{P})$ now can be reduced to maximization of a smooth function subject to a smooth constraint,

and in this kind of problem Lagrange multipliers can provide much information on the structure of the extremal graphs. For example, such approach was used successfully in [28].

We showed that for $\alpha > r$ inequality (14) follows immediately from the case $\alpha = r$, but our proof of the case $1 < \alpha \leq r$ is not easy. The proof, given in Section 4, is quite technical and mostly analytic. It is based on several lemmas of their own interest, which are presented in Section 3.

Moreover, for $r = 2$ the value of $\pi(\mathcal{P})$ can be characterized explicitly and so one can establish $\lambda^{(a)}(\mathcal{P})$ as well for all $\alpha > 1$. We give here a short proof for monotone properties, referring the reader to [31] for general hereditary properties.

Theorem 13 *Let $\alpha > 1$. If $\mathcal{P}$ is a monotone property of 2-graphs, then*

$$\lambda^{(a)}(\mathcal{P}) = \frac{r-2}{r-1},$$

where $r = \min \{\chi(G) : G \notin \mathcal{P}\}$.

It is not difficult to find a hereditary property $\mathcal{P}$ of 2-graphs for which $\lambda^{(1)}(\mathcal{P}) > \pi(\mathcal{P})$. Indeed, let $K_3(1, 2, 2)$ be the blow-up of a triangle and let $\mathcal{P} = \text{Mon}(K_3(1, 2, 2))$. First, Theorem 11 and Motzkin-Straus's result imply that for every $\alpha > 1$,

$$\lambda^{(\alpha)}(\text{Mon}(K_3(1, 2, 2))) = \lambda^{(\alpha)}(\text{Mon}(K_3)) = \lambda^{(1)}(\text{Mon}(K_3)) = 1/2.$$

Now, taking G_n to be the graph consisting of a K_4 and $n - 4$ isolated vertices, we see that $G_n \in \mathcal{P}_n$ for $n \geq 4$. But $\lambda^{(1)}(G_n) = 3/4$, and so $\lambda^{(1)}(\mathcal{P}) \geq 3/4 > 1/2$.

Obviously, Theorem 12 puts in focus hereditary properties $\mathcal{P}$ for which $\lambda^{(1)}(\mathcal{P})$ also satisfies (14); thus the definition: *a hereditary property P of r -graphs is called **flat** if $\lambda^{(1)}(\mathcal{P}) = \pi(\mathcal{P})$.*

It turns out that flat properties possess truly remarkable features with respect to extremal problems, some of which are presented in the next subsection.

1.6 Flat properties

Let us note that, in general, $\pi(\mathcal{P})$ alone is not sufficient to estimate $ex(\mathcal{P}, n)$ for small values of n and for arbitrary hereditary property $\mathcal{P}$. However, flat properties allow for tight, explicit upper bounds on $ex(\mathcal{P}, n)$ and $\lambda^{(\alpha)}(\mathcal{P})$. To emphasize the substance of the general statements in this subsection, we first outline a class of flat properties, whose study has been started by Sidorenko [37], albeit in a different setting.

A graph property P is said to be multiplicative if $G \in P_n$ implies that $G(k_1, \dots, k_n) \in P$ for every vector of positive integers $k_1, \dots, k_n$. This is to say, a multiplicative property contains the blow-ups of all its members.

Multiplicative properties are quite convenient for extremal graph theory, and they are ubiquitous as well. Indeed, following Sidorenko [37], call a graph F **covering** if every two vertices of F are contained in an edge. Clearly, complete r -graphs are covering, and for $r = 2$ these are the only covering graphs, but for $r \geq 3$ there are many noncomplete ones. For example, the Fano plane 3-graph F_7 is a noncomplete covering graph. Obviously, if F is a covering graph, then $\text{Mon}(F)$ is both a hereditary and a multiplicative property.

Below we illustrate Theorems 15 and 16 using F_7 as forbidden graph because it is covering and $\pi(\text{Mon}(F_7))$ is known. To keep our presentation focused, we stick to F_7 only, but there are other graphs with the same properties; for instance, Keevash in [15], Sec. 14, lists several such graphs, like “expanded

triangle”, “3-book with 3 pages”, “4-book with 4 pages” and others. Using these and similar references, the reader may easily come up with other illustrations. Let us point that these applications are new and are not available in the literature.

Two other examples of hereditary, multiplicative properties are based on vertex colorings. Recall that the **chromatic number** $\chi(G)$ of an r -graph G is the smallest number χ such that $V(G)$ can be partitioned into χ edgeless sets. Likewise, the **weak chromatic number** $\underline{\chi}(G)$ is the smallest number χ such that $V(G)$ can be partitioned into χ sets so that every set intersects every edge in at most one vertex.

Let now $\mathcal{C}(p)$ be the family of all r -graphs G with $\chi(G) \leq p$ and $\underline{\mathcal{C}}(q)$ be the family of all r -graphs G with $\underline{\chi}(G) \leq p$. Note first that $\mathcal{C}(p)$ and $\underline{\mathcal{C}}(q)$ are hereditary and multiplicative properties, so they are also flat. This statement is more or less obvious, but it does not follow by forbidding covering subgraphs.

The following proposition summarizes the principal facts about $\mathcal{C}(p)$ and $\underline{\mathcal{C}}(q)$.

Proposition 14 *For all p, q the classes $\mathcal{C}(p)$ and $\underline{\mathcal{C}}(q)$ are hereditary and multiplicative properties, and*

$$\pi(\mathcal{C}(p)) = (1 - p^{-r+1}) \quad \text{and} \quad \pi(\underline{\mathcal{C}}(q)) = r! \binom{q}{r} q^{-r}.$$

The motivation for the next result comes from Sidorenko [37], Theorem 2.6, who proved that if F is a covering graph, then

$$\lambda^{(1)}(\text{Mon}(F)) = \pi(\text{Mon}(F)). \quad (15)$$

This can be recast in our terminology as: *if F is a covering graph, then $\text{Mon}(F)$ is a flat property.* To analyze the underpinnings of this result, note that $\text{Mon}(F)$ is both hereditary and multiplicative property. The following theorem gives a natural generalization of (15).

Theorem 15 *If $\mathcal{P}$ is a hereditary, multiplicative property, then it is flat; that is to say,*

$$\lambda^{(\alpha)}(\mathcal{P}) = \pi(\mathcal{P}) \quad (16)$$

for every $\alpha \geq 1$.

To illustrate the usability of Theorem 15 note that the 3-graph F_7 is covering, and, as determined in [10] and [14], $\pi(\text{Mon}(F_7)) = 3/4$, so we immediately get that if $\alpha \geq 1$, then

$$\lambda^{(\alpha)}(\text{Mon}(F_7)) = 3/2.$$

However, below we show that even more convenient bounds are available in this and similar cases. Indeed a distinctive feature of all flat properties, and the one that justifies the introduction of the concept, is the fact that there exist neat and tight upper bounds on $\lambda^{(\alpha)}(G)$ and $e(G)$ for every graph G that belongs to a flat property. This claim is substantiated in Theorems 16 and 19 below.

Theorem 16 *If $\mathcal{P}$ is a flat property, and $G \in \mathcal{P}_n$, then*

$$e(G) \leq \pi(\mathcal{P}) n^r / r!. \quad (17)$$

and for every $\alpha \geq 1$,

$$\lambda^{(\alpha)}(G) \leq \pi(\mathcal{P}) n^{r-r/\alpha}. \quad (18)$$

Both inequalities (17) and (18) are tight.

When $\mathcal{P} = \text{Mon}(F)$ and F is a covering graph, the bound (17) has been proved by Sidorenko in [37], Theorem 2.3. Clearly, Theorem 16 is much more general, although its proof is similar to that of Sidorenko. Taking again the Fano plane as an example, we obtain the following tight inequality:

Corollary 17 *If G is a 3-graph of order n , not containing the Fano plane, then for all $\alpha \geq 1$,*

$$\lambda^{(\alpha)}(G) \leq \frac{3}{4} n^{3-3/\alpha}. \quad (19)$$

This inequality is essentially equivalent to Corollary 3 in [16], albeit it is somewhat less precise. We believe however, that Theorem 16 shows clearly why such a result is possible at all.

With respect to chromatic number, an early result of Cvetković [5] states: *if G is a 2-graph of order n and chromatic number χ , then*

$$\lambda(G) \leq \frac{\chi-1}{\chi} n.$$

This bound easily generalizes for hypergraphs.

Corollary 18 *Let G be an r -graph of order n and let $\alpha \geq 1$.*

(i) *If $\chi(G) = \chi$, then*

$$\lambda^{(\alpha)}(G) \leq (1 - \chi^{-r+1}) n^{r-r/\alpha};$$

(ii) *If $\underline{\chi}(G) = \underline{\chi}$, then*

$$\lambda^{(\alpha)}(G) \leq r! \binom{\underline{\chi}}{r} \underline{\chi}^{-r} n^{r-r/\alpha}.$$

Furthermore, recalling that complete graphs are the only covering 2-graphs, it becomes clear that the bound (18) is analogous to Wilf's bound [39]: *if G is a 2-graph of order n and clique number ω , then*

$$\lambda(G) \leq \frac{\omega-1}{\omega} n. \quad (20)$$

Inequality (20) has been improved by a subtler inequality in [24], namely: *if G is a 2-graph with m edges and clique number ω , then*

$$\lambda(G) \leq \sqrt{\frac{2(\omega-1)}{\omega} m}. \quad (21)$$

To see that (21) implies (20) it is enough to recall the Turán bound $m \leq (1 - 1/\omega) n^2/2$. It turns out that the proof of (21) generalizes to hypergraphs, giving the following theorem, which strengthens (18) exactly as (21) strengthens (20).

Theorem 19 *If $\mathcal{P}$ is a flat property, and $G \in \mathcal{P}$, then*

$$\lambda^{(\alpha)}(G) \leq \pi(G)^{1/\alpha} (r!e(G))^{1-1/\alpha}. \quad (22)$$

Let us emphasize the peculiar fact that the bound (22) does not depend on the order of G , but it is asymptotically tight in many cases. In particular, for 3-graphs with no F_7 we obtain the following tight bound:

Corollary 20 *If G is a 3-graph with m edges, and G does not contain the Fano plane, then*

$$\lambda^{(\alpha)}(G) \leq 3 \cdot 2^{1-3/\alpha} m^{1-1/\alpha}.$$

Finally, we shall use (22) to improve the inequality

$$\lambda^{(\alpha)}(G) \leq (r!e(G))^{1-1/\alpha},$$

given in Lemma 5 of [16]. First, note that Proposition 14, together with Theorem 19, gives the following general bounds:

Theorem 21 *If G is an r -graph and $\alpha \geq 1$, then*

$$\lambda^{(\alpha)}(G) \leq \left(1 - \chi(G)^{-r+1}\right)^{1/\alpha} (r!e(G))^{1-1/\alpha}.$$

and

$$\lambda^{(\alpha)}(G) \leq r! \binom{\underline{\chi}(G)}{r}^{1/\alpha} \underline{\chi}(G)^{-r/\alpha} e(G)^{1-1/\alpha}.$$

In particular, in view of $\chi(G) \leq v(G)/(r-1)$ and $\underline{\chi}(G) \leq v(G)$, we obtain simple bounds in terms of the order and size:

Corollary 22 *If $\alpha \geq 1$ and G is an r -graph of order n , then*

$$\lambda^{(\alpha)}(G) \leq \left(1 - \left(\frac{n}{r-1}\right)^{-r+1}\right)^{1/\alpha} (r!e(G))^{1-1/\alpha}.$$

and

$$\lambda^{(\alpha)}(G) \leq r! \binom{n}{r}^{1/\alpha} n^{-r/\alpha} e(G)^{1-1/\alpha}$$

If G contains at least one edge, then

$$\lambda^{(\alpha)}(G) < (r!e(G))^{1-1/\alpha}.$$

The above results lead to the natural question: *Is there a flat property that is not multiplicative?* The answer is yes, there exists a flat property $\mathcal{P}$ of 2-graphs that is not multiplicative. Indeed, let $\mathcal{P} = \text{Her}(C_4)$, that is to say, $\mathcal{P}$ is the class of all graphs with no induced 4-cycle. Trivially, all complete graphs belong to $\mathcal{P}$ and so

$$\lambda^{(\alpha)}(\mathcal{P}) = \pi(G) = 1 \quad \text{for all } \alpha \geq 1.$$

However, obviously $\mathcal{P}$ is not multiplicative, as $C_4 = K_2(2, 2)$.

Analyzing the above example, we come up with the following sufficient condition for flat properties.

Theorem 23 *Let $\mathcal{F}$ be a set of r -graphs each of which is a blow-up of a covering graph. Then $\text{Her}(\mathcal{F})$ is flat.*

Apparently Theorem 23 greatly extends the range of flat properties, however further work is needed to determine the limits of its applicability.

The rest of the paper is organized as follows: in Section 2 we give general results for the parameters $\lambda^{(\alpha)}(G)$ and in particular for $\lambda(G)$. In Section 3 we prove two useful lemmas which are needed in the proof of Theorem 15, but are of separate interest as well. The proofs of the various statements above are presented in Section 4. The paper ends with a summary discussion and open problems in Section 5.

2 Some properties of $\lambda(G)$ and $\lambda^{(\alpha)}(G)$

Below we write $\binom{X}{k}$ for the set of all k -subsets of a set X . Let us recall also the notation $(n)_r$ for the falling factorial $n!/(n-k)!$

$$(n)_r = \frac{n!}{(n-k)!} = n(n-1) \cdots (n-k+1).$$

In our proofs we shall use extensively Jensen's and Maclauren's inequalities; the reader is referred to [12] for ground material.

Most of the basic results about $\lambda^{(\alpha)}(G)$ appear here for the first time, and we hope that they will be useful to other researchers.

Let G be an r -graph of order n . Since its polyform $P_G(\mathbf{x})$ is homogenous, we find that

$$\lambda(G) = \max_{\|\mathbf{x}\|_r=1} P_G(\mathbf{x}) = \max_{\mathbf{x} \neq 0} \frac{P_G(\mathbf{x})}{\|\mathbf{x}\|_r^r} = \max_{\mathbf{x} \neq 0} P_G\left(\frac{1}{\|\mathbf{x}\|_r} \mathbf{x}\right). \quad (23)$$

We shall call a nonzero real vector $\mathbf{x} = (x_1, \dots, x_n)$ an **eigenvector** to $\lambda(G)$ if

$$\lambda(G) = P_G\left(\frac{1}{\|\mathbf{x}\|_r} \mathbf{x}\right).$$

Note that relation (23) holds for $\lambda(G)$, but it is not true for $\lambda^{(\alpha)}(G)$ if $\alpha \neq r$. This fact corroborates again the exclusivity of $\lambda(G)$. However, let us note a useful inequality for the general $\lambda^{(\alpha)}(G)$, which we shall use later with no explicit reference.

Proposition 24 *Let $\alpha \geq 1$. If G is an r -graph of order n , and $\mathbf{x} = (x_1, \dots, x_n)$ is any real vector, then*

$$P_G(\mathbf{x}) \leq \lambda^{(\alpha)}(G) \|\mathbf{x}\|_\alpha^r.$$

Here are two other obvious facts:

Proposition 25 *If G is an r -graph with at least one edge then $\lambda^{(\alpha)}(G) \geq r!r^{-r/\alpha}$. If H is a subgraph of G , then $\lambda^{(\alpha)}(H) \leq \lambda^{(\alpha)}(G)$.*

We state below a handy fact that will be used later with no explicit reference.

Proposition 26 *For any $\alpha \geq 1$ there is a nonnegative vector $\mathbf{x}$ such that $\|\mathbf{x}\|_\alpha = 1$ and $\lambda^{(\alpha)}(G) = P_G(\mathbf{x})$.*

Indeed, if $\mathbf{x} = (x_1, \dots, x_n)$ is a vector such that $\|\mathbf{x}\|_\alpha = 1$ and

$$\lambda^{(\alpha)}(G) = P_G(\mathbf{x}) = \max_{\|\mathbf{x}\|_\alpha=1} P_G(\mathbf{x}), \quad (24)$$

then $\lambda^{(\alpha)}(G) = P_G(|x_1|, \dots, |x_n|)$.

If $\alpha = r$, there are stronger statements, which are analogous to statements in the Perron-Frobenius theory for nonnegative matrices. The following crucial statement can be deduced from the results of Friedland, Gauber and Han [9], but an independent, direct proof has been given also by Cooper and Dutle in [4].

Theorem 27 *Let G be a connected r -graph. Then $\lambda(A)$ has a positive eigenvector, which is unique up to scaling.* $\square$

Note, however, that for $r \geq 3$, even if G is a connected r -graph, $\lambda(G)$ may have eigenvectors with negative and positive entries; for example, if G is a one edge graph, then the r -vector $(1, 1, \dots, 1)$ is an eigenvector to $\lambda(G)$, but so is the vector $(-1, -1, 1, \dots, 1)$ as well.

Let now $\alpha > 1$ and let $\mathbf{x} = (x_1, \dots, x_n)$ be a nonnegative vector satisfying $\|\mathbf{x}\|_\alpha = 1$ and (24). Using Lagrange multipliers, we find that there exists μ such that, for every $k = 1, \dots, n$,

$$\mu \alpha x_k^{\alpha-1} = (r-1)! \sum_{\{k, i_1, \dots, i_{r-1}\} \in E(G)} x_{i_1} \cdots x_{i_{r-1}}.$$

Multiplying the k 'th equation by x_k and adding them all, we find that $\mu \alpha = \lambda^{(\alpha)}(G)$, and so, the numbers $x_1, \dots, x_n$ satisfy the n equations

$$\lambda^{(\alpha)}(G) x_k^{\alpha-1} = (r-1)! \sum_{\{k, i_1, \dots, i_{r-1}\} \in E(G)} x_{i_1} \cdots x_{i_{r-1}}, \quad 1 \leq k \leq n. \quad (25)$$

For $\alpha > 1$ these equations are a powerful tool in the study of $\lambda^{(\alpha)}(G)$ and particularly of $\lambda(G)$, but they are not always available for $\alpha = 1$.

Next, we shall prove a relation between $\lambda^{(\alpha)}(G)$ and $\lambda^{(\alpha)}(G(k, \dots, k))$, where $G(k, \dots, k)$ is a uniform blow-up of G .

Proposition 28 *If G is an r -graph and $k \geq 1$ is an integer, then*

$$\lambda^{(\alpha)}(G(k, \dots, k)) = k^{r-r/\alpha} \lambda^{(\alpha)}(G).$$

Proof By definition, the vertex set $V(G(k, \dots, k))$ of $G(k, \dots, k)$ can be partitioned into n disjoint sets $U_1, \dots, U_n$ each consisting of k vertices. Also, if $\{i_1, \dots, i_r\} \in E(G)$, then $\{j_1, \dots, j_r\} \in E(G(k, \dots, k))$ for every $j_1 \in U_{i_1}, j_2 \in U_{i_2}, \dots, j_r \in U_{i_r}$. We shall prove first that

$$\lambda^{(\alpha)}(G(k, \dots, k)) \geq k^{r-r/\alpha} \lambda^{(\alpha)}(G). \quad (26)$$

Let $\mathbf{x} = (x_1, \dots, x_n)$ be a nonnegative vector such that $\|\mathbf{x}\|_\alpha = 1$ and $\lambda^{(\alpha)}(G) = P_G(\mathbf{x})$. For every $i \in [n]$ and every $j \in U_i$, set

$$y_j = \frac{1}{k^{1/\alpha}} x_i$$

The vector $\mathbf{y} = (y_1, \dots, y_{nk})$ satisfies $\|\mathbf{y}\|_\alpha = 1$, and therefore,

$$\lambda^{(\alpha)}(G(k, \dots, k)) \geq P_{G(k, \dots, k)}(\mathbf{y}) = \frac{1}{k^{r/\alpha}} k^r P_G(\mathbf{x}) = k^{r-r/\alpha} \lambda^{(\alpha)}(G),$$

proving (26). To complete the proof of the proposition, we shall show that

$$\lambda^{(\alpha)}(G(k, \dots, k)) \leq k^{r-r/\alpha} \lambda^{(\alpha)}(G). \quad (27)$$

Let $\mathbf{x} = (x_1, \dots, x_{nk})$ be a nonnegative kn -vector such that $\|\mathbf{x}\|_\alpha = 1$ and

$$\lambda^{(\alpha)}(G(k, \dots, k)) = P_{G(k, \dots, k)}(\mathbf{x}).$$

By definition,

$$\begin{aligned}
P_{G(k,\dots,k)}(\mathbf{x}) &= r! \sum_{\{i_1,\dots,i_r\} \in E(G(k,\dots,k))} x_{i_1} x_{i_2} \cdots x_{i_r} \\
&= r! \sum_{\{i_1,\dots,i_r\} \in E(G)} \left(\sum_{j \in U_{i_1}} x_j \right) \left(\sum_{j \in U_{i_2}} x_j \right) \cdots \left(\sum_{j \in U_{i_r}} x_j \right)
\end{aligned}$$

Next, for every $s \in [n]$, using Jensen's inequality, we see that

$$\sum_{j \in U_s} x_j \leq k^{1-1/\alpha} \left(\sum_{j \in U_s} x_j^\alpha \right)^{1/\alpha}.$$

Now, setting for every $s \in [n]$,

$$y_s = \left(\sum_{j \in U_s} x_j^\alpha \right)^{1/\alpha},$$

we find a vector $\mathbf{y} = (y_1, \dots, y_n)$ with $\|\mathbf{y}\|_\alpha = 1$. Also,

$$\begin{aligned}
\lambda^{(\alpha)}(G(k, \dots, k)) &= P_{G(k,\dots,k)}(\mathbf{x}) \\
&= r! \sum_{\{i_1,\dots,i_r\} \in E(G)} \left(\sum_{j \in U_{i_1}} x_j \right) \left(\sum_{j \in U_{i_2}} x_j \right) \cdots \left(\sum_{j \in U_{i_r}} x_j \right) \\
&\leq r! k^{r-r/\alpha} \sum_{\{i_1,\dots,i_r\} \in E(G)} y_{i_1} y_{i_2} \cdots y_{i_r} = k^{r-r/\alpha} P_G(\mathbf{y}) \\
&\leq k^{r-r/\alpha} \lambda^{(\alpha)}(G).
\end{aligned}$$

This completes the proof of (27), and with (26), also the proof of Proposition 28. $\square$

Finally, we give a perturbation bound on $\lambda^{(\alpha)}(G)$, which is used to estimate how much changes $\lambda^{(\alpha)}(G)$, when edges of G are changed.

Proposition 29 *Let $\alpha \leq 1$, $k \geq 1$ and G_1 and G_2 be r -graphs on the same vertex set. If G_1 and G_2 differ in at most k edges, then*

$$\left| \lambda^{(\alpha)}(G_1) - \lambda^{(\alpha)}(G_2) \right| \leq (r!k)^{1-1/\alpha}.$$

Proof Let $V = V(G_1) = V(G_2)$ and write G_{12} for the r -graph with $V(G_{12}) = V$ and $E(G_{12}) = E(G_1) \cap E(G_2)$. We may and shall assume that $\lambda^{(\alpha)}(G_1) \geq \lambda^{(\alpha)}(G_2)$. Write G_3 for the r -graph with $V(G_3) = V$ and $E(G_3) = E(G_1) \setminus E(G_2)$. In view of $G_{12} \subset G_2$, we have

$$\begin{aligned}
0 &\leq \lambda^{(\alpha)}(G_1) - \lambda^{(\alpha)}(G_2) = \lambda^{(\alpha)}(G_1) - \lambda^{(\alpha)}(G_{12}) - \left(\lambda^{(\alpha)}(G_2) - \lambda^{(\alpha)}(G_{12}) \right) \\
&\leq \lambda^{(\alpha)}(G_1) - \lambda^{(\alpha)}(G_{12}) = \lambda^{(\alpha)}(G_3) \leq (r!e(G_3))^{1-1/\alpha} \\
&\leq (r!k)^{1-1/\alpha},
\end{aligned}$$

proving Proposition 29. $\square$

3 Two lemmas about critical vectors to $\lambda^{(\alpha)}(G)$

A useful result in spectral extremal theory for 2-graphs is the following bound from [26]:

Let G be an 2-graph with minimum degree δ , and let $x = (x_1, \dots, x_n)$ be a nonnegative eigenvector to $\lambda(G)$ with $\|x\|_2 = 1$. Then the value $x = \min\{x_1, \dots, x_n\}$ satisfies

$$x^2 \left(\lambda(G)^2 + \delta n - \delta^2 \right) \leq \delta \quad (28)$$

The bound (28) is exact for many different graphs, and as explained in [27], it has been crucial in proving upper bounds on $\lambda(G)$ by induction on the number of vertices of G . Very likely, a similar bound for hypergraphs would be useful as well. Below we state and prove such a result; despite its awkward form, for $r = \alpha = 2$ it yields precisely (28); moreover, it is crucial for the proof of Theorem 12.

Lemma 30 *Let $1 \leq \alpha \leq r$, and let G be an r -graph of order n , with minimum degree δ and with $\lambda^{(\alpha)}(G) = \lambda$. Let $\mathbf{x} = (x_1, \dots, x_n)$ be a nonnegative vector such that $\|\mathbf{x}\|_\alpha = 1$ and $\lambda = P_G(\mathbf{x})$. Then the value $x = \min\{x_1, \dots, x_n\}$ satisfies*

$$\left(\left(\frac{\lambda n^{r/\alpha-1}}{(r-1)!} \right)^\alpha - \delta^\alpha \right) x^{\alpha(r-1)} \leq \binom{n-1}{r-1} \delta^{\alpha-1} \left(\frac{(1-x^\alpha)^{r-1}}{(n-1)^{r-1}} - x^{\alpha(r-1)} \right).$$

Proof Set for short $V = V(G)$ and let $k \in V$ be a vertex of degree δ . The equation (25) for the vertex k implies that

$$\lambda x^{\alpha-1} \leq \lambda x_k^{\alpha-1} = (r-1)! \sum_{\{k, i_1, \dots, i_{r-1}\} \in E(G)} x_{i_1} \dots x_{i_{r-1}}.$$

Now, dividing by $(r-1)!$ and applying Jensen's inequality to the right-hand side, we find that

$$\left(\frac{\lambda x^{\alpha-1}}{(r-1)!} \right)^\alpha \leq \delta^{\alpha-1} \sum_{\{k, i_1, \dots, i_{r-1}\} \in E(G)} x_{i_1}^\alpha \dots x_{i_{r-1}}^\alpha. \quad (29)$$

Our next goal is to bound the quantity $\sum_{\{k, i_1, \dots, i_{r-1}\} \in E(G)} x_{i_1}^\alpha \dots x_{i_{r-1}}^\alpha$ from above. First, note that

$$\begin{aligned} & \sum_{\{k, i_1, \dots, i_{r-1}\} \in E(G)} x_{i_1}^\alpha \dots x_{i_{r-1}}^\alpha \\ &= \sum_{\{i_1, \dots, i_{r-1}\} \in \binom{V \setminus \{v_k\}}{r-1}} x_{i_1}^\alpha \dots x_{i_{r-1}}^\alpha - \left(\sum_{\{i_1, \dots, i_{r-1}\} \in \binom{V \setminus \{v_k\}}{r-1} \text{ and } \{k, i_1, \dots, i_{r-1}\} \notin E(G)} x_{i_1}^\alpha \dots x_{i_{r-1}}^\alpha \right) \\ &\leq \sum_{\{i_1, \dots, i_{r-1}\} \in \binom{V \setminus \{v_k\}}{r-1}} x_{i_1}^\alpha \dots x_{i_{r-1}}^\alpha - \left(\sum_{\{i_1, \dots, i_{r-1}\} \in \binom{V \setminus \{v_k\}}{r-1} \text{ and } \{k, i_1, \dots, i_{r-1}\} \notin E(G)} x^{\alpha(r-1)} \right) \\ &= \sum_{\{i_1, \dots, i_{r-1}\} \in \binom{V \setminus \{v_k\}}{r-1}} x_{i_1}^\alpha \dots x_{i_{r-1}}^\alpha - \left(\binom{n-1}{r-1} - \delta \right) x^{\alpha(r-1)}. \end{aligned} \quad (30)$$

Next, applying Maclauren's inequality for the $(r-1)$ 'th symmetric function of the variables $x_i^\alpha, i \in V \setminus \{v_k\}$, we find that

$$\begin{aligned} \frac{1}{\binom{n-1}{r-1}} \sum_{\{i_1, \dots, i_{r-1}\} \in \binom{V \setminus \{v_k\}}{r-1}} x_{i_1}^\alpha \dots x_{i_{r-1}}^\alpha &\leq \left(\frac{1}{n-1} \sum_{i \in V \setminus \{k\}} x_i^\alpha \right)^{r-1} = \frac{1}{(n-1)^{r-1}} (1 - x_k^\alpha)^{r-1} \\ &\leq \frac{1}{(n-1)^{r-1}} (1 - x^\alpha)^{r-1}. \end{aligned}$$

Hence, replacing in (30), we obtain the desired bound

$$\sum_{\{k, i_1, \dots, i_{r-1}\} \in E(G)} x_{i_1}^\alpha \dots x_{i_{r-1}}^\alpha \leq \frac{\binom{n-1}{r-1}}{(n-1)^{r-1}} (1 - x^\alpha)^{r-1} - \left(\binom{n-1}{r-1} - \delta \right) x^{\alpha(r-1)}.$$

Returning back to (29), we see that

$$\left(\frac{\lambda}{(r-1)!} \right)^\alpha x^{(\alpha-1)\alpha} \leq \binom{n-1}{r-1} \delta^{\alpha-1} \left(\frac{(1-x^\alpha)^{r-1}}{(n-1)^{r-1}} - x^{\alpha(r-1)} \right) + \delta^\alpha x^{\alpha(r-1)}.$$

Since $\alpha \leq r$ and $x \leq n^{-1/\alpha}$, we see that $x^{\alpha-r} \geq n^{-(\alpha-r)/\alpha} = n^{r/\alpha-1}$ and therefore,

$$\frac{\lambda n^{r/\alpha-1}}{(r-1)!} x^{r-1} \leq \frac{\lambda}{(r-1)!} x^{\alpha-1}.$$

Hence,

$$\begin{aligned} \left(\frac{\lambda n^{r/\alpha-1}}{(r-1)!} \right)^\alpha x^{\alpha(r-1)} &\leq \left(\frac{\lambda}{(r-1)!} \right)^\alpha x^{(\alpha-1)\alpha} \\ &\leq \binom{n-1}{r-1} \delta^{\alpha-1} \left(\frac{(1-x^\alpha)^{r-1}}{(n-1)^{r-1}} - x^{\alpha(r-1)} \right) + \delta^\alpha x^{\alpha(r-1)}, \end{aligned}$$

and so,

$$\left(\left(\frac{\lambda n^{r/\alpha-1}}{(r-1)!} \right)^\alpha - \delta^\alpha \right) x^{\alpha(r-1)} \leq \binom{n-1}{r-1} \delta^{\alpha-1} \left(\frac{(1-x^\alpha)^{r-1}}{(n-1)^{r-1}} - x^{\alpha(r-1)} \right),$$

completing the proof of Lemma 30. □

If G is an r -graph of order n , with minimum degree δ , from (2) we see that

$$\frac{\lambda^{(\alpha)}(G) n^{r/\alpha-1}}{(r-1)!} \geq \frac{re(G)}{n} \geq \delta. \quad (31)$$

In the proof of Lemma 32, we shall need the following simple consequence of this fact.

Proposition 31 *If $\alpha \geq 1$, $r \geq 2$, and G is an r -graph of order n , with minimum degree δ , then*

$$\left(\frac{\lambda^{(\alpha)}(G) n^{r/\alpha-1}}{(r-1)!} \right)^\alpha - \delta^\alpha \geq \left(\frac{\lambda^{(\alpha)}(G) n^{r/\alpha-1}}{(r-1)!} - \delta \right) \alpha \delta^{\alpha-1} \quad (32)$$

Proof Indeed, setting for short

$$\frac{\lambda^{(\alpha)}(G) n^{r/\alpha-1}}{(r-1)!} = b,$$

inequality (32) is equivalent to

$$b^\alpha - \delta^\alpha \geq (b - \delta) \alpha \delta^{\alpha-1}.$$

But (31) implies that $b \geq \delta$, and using Bernoulli's inequality, we find that

$$b^\alpha - \delta^\alpha = \delta^\alpha \left(\frac{b^\alpha}{\delta^\alpha} - 1 \right) = \delta^\alpha \left(\left(1 + \frac{b - \delta}{\delta} \right)^\alpha - 1 \right) \geq \delta^\alpha \left(\frac{\alpha (b - \delta)}{\delta} \right) = (b - \delta) \alpha \delta^{\alpha-1},$$

completing the proof of the proposition. $\square$

Recall that inequality (31) is obtained by taking a vector $(x_1, \dots, x_n)$ with $x_1 = \dots = x_n = n^{-1/\alpha}$. Now let $\mathbf{x} = (x_1, \dots, x_n)$ be a nonnegative vector such that $\|\mathbf{x}\|_\alpha = 1$ and $\lambda^{(\alpha)}(G) = P_G(\mathbf{x})$. It turns out that if the entries of $\mathbf{x}$ are close to $n^{-1/\alpha}$, then the bound (31) can be inverted to some extent, which also implies that the graph is almost regular. The following technical lemma gives a quantitative form of this statement.

Lemma 32 *Let $1 < \alpha \leq r$, let G be an r -graph of sufficiently large order n , with minimum degree δ and with $\lambda^{(\alpha)}(G) = \lambda$. Let $\mathbf{x} = (x_1, \dots, x_n)$ be a nonnegative vector such that $\|\mathbf{x}\|_\alpha = 1$ and $\lambda = P_G(\mathbf{x})$. If the value $x = \min\{x_1, \dots, x_n\}$ satisfies*

$$x^\alpha \geq \frac{1}{n} \left(1 - \frac{1}{(\alpha - 1) \log n} \right), \quad (33)$$

then

$$\frac{\lambda n^{r/\alpha-1}}{(r-1)!} \leq \delta + \frac{2r}{\alpha(\alpha-1) \log n} \binom{n-1}{r-1}.$$

Proof To begin with, Lemma 30 gives

$$\left(\left(\frac{\lambda n^{r/\alpha-1}}{(r-1)!} \right)^\alpha - \delta^\alpha \right) x^{\alpha(r-1)} \leq \binom{n-1}{r-1} \delta^{\alpha-1} \left(\frac{(1-x^\alpha)^{r-1}}{(n-1)^{r-1}} - x^{\alpha(r-1)} \right).$$

Now, since the premise (33) implies that $x > 0$, we can rearrange the above inequality to

$$\left(\frac{\lambda n^{r/\alpha-1}}{(r-1)!} \right)^\alpha - \delta^\alpha \leq \binom{n-1}{r-1} \delta^{\alpha-1} \left(\frac{1}{(n-1)^{r-1}} \frac{(1-x^\alpha)^{r-1}}{x^{\alpha(r-1)}} - 1 \right). \quad (34)$$

Obviously, the expression $(1-y)/y$ decreases with y ; in particular, the premise

$$x^\alpha \geq \frac{1}{n} \left(1 - \frac{1}{(\alpha - 1) \log n} \right)$$

implies that

$$\begin{aligned}
\frac{1-x^\alpha}{x^\alpha} &\leq \frac{n-1 + \frac{1}{(\alpha-1)\log n}}{1 - \frac{1}{(\alpha-1)\log n}} = \frac{(n-1)(\alpha-1)\log n + 1}{(\alpha-1)\log n - 1} \\
&= (n-1) \frac{\log n + \frac{1}{(n-1)(\alpha-1)}}{\log n - \frac{1}{(\alpha-1)}} \\
&= (n-1) \left(1 + \frac{n}{(\alpha-1)(n-1)\left(\log n - \frac{1}{(\alpha-1)}\right)} \right).
\end{aligned}$$

Now, bounding the expression $(1-x^\alpha)/x^\alpha$ in (34), we get

$$\left(\frac{\lambda n^{r/\alpha-1}}{((r-1)!)} \right)^\alpha - \delta^\alpha < \binom{n-1}{r-1} \delta^{\alpha-1} \left(\left(1 + \frac{n}{(\alpha-1)(n-1)\left(\log n - \frac{1}{(\alpha-1)}\right)} \right)^{r-1} - 1 \right). \quad (35)$$

Next, assuming that n is large enough and using Bernoulli's inequality, we find that

$$\begin{aligned}
\left(1 + \frac{1}{(\alpha-1)(1-1/n)\left(\log n - \frac{1}{(\alpha-1)}\right)} \right)^{r-1} &\leq 1 + \frac{r-1}{(\alpha-1)(1-1/n)\left(\log n - \frac{1}{(\alpha-1)}\right) - r} \\
&\leq 1 + \frac{2r}{(\alpha-1)\log n}.
\end{aligned}$$

Replacing this bound in (35), we get

$$\left(\frac{\lambda n^{r/\alpha-1}}{((r-1)!)} \right)^\alpha - \delta^\alpha < \binom{n-1}{r-1} \delta^{\alpha-1} \frac{2r}{(\alpha-1)\log n}. \quad (36)$$

On the other hand, (32) implies

$$\left(\frac{\lambda n^{r/\alpha-1}}{(r-1)!} - \delta \right) \alpha \delta^{\alpha-1} \leq \binom{n-1}{r-1} \delta^{\alpha-1} \frac{2r}{(\alpha-1)\log n},$$

and so,

$$\frac{\lambda n^{r/\alpha-1}}{(r-1)!} \leq \delta + \frac{2r}{\alpha(\alpha-1)\log n} \binom{n-1}{r-1},$$

completing the proof of Lemma 32. □

4 Proofs

The proofs given below follow the order of appearance of the statements in Section 1, except for the proof of Theorem 13, given at the end of the section.

Proof of Proposition 3 Let $\mathbf{x} = (x_1, \dots, x_n)$ be a nonnegative vector such that $\|\mathbf{x}\|_\alpha = 1$ and $\lambda^{(\alpha)}(G) = P_G(\mathbf{x})$. Assume that $\alpha \geq r$, let $x = \max\{x_1, \dots, x_n\}$, and let $k \in V(G)$ be a vertex for which $x_k = x$. From equations (25) we have

$$\frac{\lambda^{(\alpha)}(G)}{(r-1)!} x^{\alpha-1} = \sum_{\{k, i_1, \dots, i_{r-1}\} \in E(G)} x_{i_1} \cdots x_{i_{r-1}} \leq \Delta x^{r-1}$$

Since $x \geq n^{-1/\alpha}$ and $\alpha \geq r$, we find that

$$\frac{\lambda^{(\alpha)}(G)}{(r-1)!} \leq \Delta x^{r-\alpha} \leq \Delta \left(n^{-1/\alpha}\right)^{r-\alpha} = \frac{\Delta}{n^{r/\alpha-1}},$$

proving (3). Now if we have equality in the above, then $x = n^{-1/\alpha}$ and so $x_1 = \dots = x_n = n^{-1/\alpha}$. Thus, equations (25) show that all degrees are equal to $\lambda^{(\alpha)}(G) / (r-1)! = \Delta$, and G is regular. On the other hand,

$$\frac{re(G)}{n^{r/\alpha}} \leq \frac{\lambda^{(\alpha)}(G)}{(r-1)!} \leq \frac{\Delta}{n^{r/\alpha-1}},$$

and so if G is regular, then $\lambda^{(\alpha)}(G) / (r-1)! = \Delta n^{1-r/\alpha}$, completing the proof of (i).

To prove (ii), assume that $1 \leq \alpha < r$ and let G be the union of n disjoint complete r -graph on $k > r$ vertices. It is easy to see that

$$\lambda^{(\alpha)}(G_0) = P_{G_0}\left(\left(k^{-1/\alpha}, \dots, k^{-1/\alpha}\right)\right) = r! \binom{k}{r} k^{-r/\alpha}$$

and $\Delta = \binom{k-1}{r-1}$. Hence,

$$\frac{\lambda^{(\alpha)}(G_0)}{(r-1)!} = \frac{\Delta}{k^{r/\alpha-1}},$$

and so (3) fails if n is large, because $r > \alpha$. This completes the proof of (ii). $\square$

There is nothing to prove in (iii), in view of $\lambda^{(\alpha)}(G) < \lambda^{(r)}(G) \leq (r-1)!\Delta$. $\square$

Proof of Proposition 4 If $\lambda^{(\alpha)}(G) = r!e(G)/n^{r/\alpha}$, then for the n -vector $\mathbf{x} = (n^{-1/\alpha}, \dots, n^{-1/\alpha})$ we see that $\lambda^{(\alpha)}(G) = P_G(\mathbf{x})$, and so $\mathbf{x}$ satisfies equations (25), which implies that all degrees are equal. If $\alpha \geq r$ and G is regular, then from

$$\frac{re(G)}{n^{r/\alpha}} \leq \frac{\lambda^{(\alpha)}(G)}{(r-1)!} \leq \frac{\Delta}{n^{r/\alpha-1}},$$

we see that $\lambda^{(\alpha)}(G) = r!e(G)/n^{r/\alpha}$. Finally, let $1 \leq \alpha < r$, fix an integer $k > r$, and take the union of n complete r -graphs of order k . As in the proof of Proposition 3 we see that

$$\lambda^{(\alpha)}(G) = r! \binom{k}{r} k^{-r/\alpha} > r!n \binom{k}{r} / n^{r/\alpha} = r!e(G) / n^{r/\alpha},$$

completing the proof of Proposition 4. $\square$

Proof of Proposition 5 Let $1 \leq \alpha < \beta$, and let $\mathbf{x} = (x_1, \dots, x_n)$ be a nonnegative vector satisfying $\|\mathbf{x}\|_\beta = 1$ and $\lambda^{(\beta)}(G) = P_G(\mathbf{x})$. Using Jensen's inequality, we see that

$$\|\mathbf{x}\|_\alpha^r = (x_1^\alpha + \dots + x_n^\alpha)^{r/\alpha} \leq \left(x_1^\beta + \dots + x_n^\beta\right)^{r/\beta} n^{r/\alpha-r/\beta} = n^{r/\alpha-r/\beta}. \quad (37)$$

Therefore,

$$\lambda^{(\beta)}(G) = P_G(\mathbf{x}) \leq \lambda^{(\alpha)}(G) \|\mathbf{x}\|_\alpha^r \leq \lambda^{(\alpha)}(G) n^{r/\alpha - r/\beta},$$

implying that $h_G(\alpha)$ is nonincreasing.

If G is non-regular, then some of the entries $x_1, \dots, x_n$ are distinct and so Jensen's inequality implies strict inequality in (37), implying in turn that

$$\lambda^{(\beta)}(G) < \lambda^{(\alpha)}(G) n^{r/\alpha - r/\beta},$$

proving Proposition 5. □

Proof of Proposition 6 Let $1 \leq \alpha < \beta$. Set for short $m = e(G)$ and let $\mathbf{x} = (x_1, \dots, x_n)$ be a nonnegative vector satisfying $\|\mathbf{x}\|_\beta = 1$ and $\lambda^{(\beta)}(G) = P_G(\mathbf{x})$. Using Jensen's inequality, we see that

$$\frac{\lambda^{(\beta)}(G)}{r!m} = \frac{1}{m} \sum_{\{i_1, \dots, i_r\} \in E(G)} x_{i_1} \dots x_{i_r} \leq \left(\frac{1}{m} \sum_{\{i_1, \dots, i_r\} \in E(G)} x_{i_1}^{\beta/\alpha} \dots x_{i_r}^{\beta/\alpha} \right)^{\alpha/\beta}.$$

Note that

$$\left(x_1^{\beta/\alpha}\right)^\alpha + \dots + \left(x_n^{\beta/\alpha}\right)^\alpha = x_1^\beta + \dots + x_n^\beta = 1,$$

and so, for the vector $\mathbf{y} = (x_1^{\beta/\alpha}, \dots, x_n^{\beta/\alpha})$ we have $\|\mathbf{y}\|_\alpha = 1$. Hence,

$$\frac{1}{m} \sum_{\{i_1, \dots, i_r\} \in E(G)} x_{i_1}^{\beta/\alpha} \dots x_{i_r}^{\beta/\alpha} = \frac{1}{r!m} P_G(\mathbf{y}) \leq \frac{1}{r!m} \lambda^{(\alpha)}(G),$$

and so,

$$\left(\frac{\lambda^{(\beta)}(G)}{r!m} \right)^\beta \leq \left(\frac{\lambda^{(\alpha)}(G)}{r!m} \right)^\alpha,$$

proving Proposition 6. □

Proof of Theorem 8 For every integer $n \geq 2$, set for short $\lambda_n^{(\alpha)} = \lambda^{(\alpha)}(\mathcal{P}, n)$. Let $G \in \mathcal{P}_n$ be such that $\lambda^{(\alpha)}(G) = \lambda_n^{(\alpha)}$ and let $\mathbf{x} = (x_1, \dots, x_n)$ be a nonnegative vector such that $\|\mathbf{x}\|_\alpha = 1$ and

$$\lambda_n^{(\alpha)} = \lambda^{(\alpha)}(G) = P_G(\mathbf{x}).$$

If $\alpha = 1$, we obviously have $\lambda_n^{(1)} \geq \lambda_{n-1}^{(1)}$. and in view of

$$\lambda_n^{(1)} = P_G(\mathbf{x}) \leq r! \sum_{1 \leq i_1 < \dots < i_r \leq n} x_{i_1} \dots x_{i_r} \leq (x_1 + \dots + x_n)^r = 1,$$

we see that the sequence $\left\{ \lambda_n^{(1)} \right\}_{n=1}^\infty$ is converging to some λ . Then,

$$\lambda = \lim_{n \rightarrow \infty} \lambda_n^{(1)} n^{r-r} = \lambda^{(1)}(\mathcal{P}),$$

proving (7)

Suppose now that $\alpha > 1$. Obviously there exists a vertex k of G such that $x_k^\alpha \leq 1/n$. Write $G - k$ for the r -graph obtained from $\mathbf{x}$ by omitting the vertex k , and let $\mathbf{x}'$ be the $(n-1)$ -vector obtained from $\mathbf{x}$ by omitting the entry x_k . Then,

$$\begin{aligned} P_{G-k}(\mathbf{x}') &= P_G(\mathbf{x}) - r!x_k \sum_{\{k, i_1, \dots, i_{r-1}\} \in E(G)} x_{i_1} \dots x_{i_{r-1}} = \\ &= \lambda^{(\alpha)}(G) - rx_k \left(\lambda^{(\alpha)}(G) x_k^{\alpha-1} \right) = \lambda_n^{(\alpha)} (1 - rx_k^\alpha) \end{aligned}$$

On the other hand, $\mathcal{P}$ is a hereditary property, so $G - k \in \mathcal{P}_{n-1}$, and therefore,

$$P_{G-k}(\mathbf{x}') \leq \lambda^{(\alpha)}(G - k) \|\mathbf{x}'\|_\alpha^r = \lambda^{(\alpha)}(G - k) (1 - x_k^\alpha)^{r/\alpha} \leq \lambda_{n-1}^{(\alpha)} (1 - x_k^\alpha)^{r/\alpha}.$$

Thus, we obtain

$$\lambda_n^{(\alpha)} \leq \lambda_{n-1}^{(\alpha)} \frac{(1 - x_k^\alpha)^{r/\alpha}}{(1 - rx_k^\alpha)}. \quad (38)$$

Note that the function

$$f(x) = \frac{(1-x)^{r/\alpha}}{1-rx}$$

is nondecreasing in x for $0 \leq x \leq 1/n$ and n sufficiently large. Indeed,

$$\begin{aligned} \frac{df(x)}{dx} &= \frac{-\frac{r}{\alpha}(1-x)^{r/\alpha-1}(1-rx) + rx(1-x)^{r/\alpha}}{(1-rx)^2} \\ &= \left(-\frac{1}{\alpha}(1-rx) + (1-x) \right) \frac{r(1-x)^{r/\alpha-1}}{(1-rx)^2} \\ &= \left(-\left(\frac{1}{\alpha} - 1 \right) + \left(\frac{r}{\alpha} - 1 \right) x \right) \frac{r(1-x)^{r/\alpha-1}}{(1-rx)^2} \geq 0 \end{aligned}$$

Here we use the fact that $1/\alpha - 1 > 0$ and that $(r/\alpha - 1)x$ tends to 0 when $n \rightarrow \infty$.

Hence, in view of (38), we find that for n large enough,

$$\lambda_n^{(\alpha)} \leq \lambda_{n-1}^{(\alpha)} f(x_k^\alpha) \leq \lambda_{n-1}^{(\alpha)} f\left(\frac{1}{n}\right) = \lambda_{n-1}^{(\alpha)} \frac{n(1-1/n)^{r/\alpha}}{(n-r)},$$

and so, (38) implies that

$$\frac{\lambda_n^{(\alpha)} n^{r/\alpha-1}}{(n-1)(n-2) \dots (n-r+1)} \leq \frac{\lambda_n^{(\alpha)} (n-1)^{r/\alpha-1}}{(n-2)(n-2) \dots (n-r)}.$$

Therefore, the sequence

$$\left\{ \frac{\lambda_n^{(\alpha)} n^{r/\alpha-1}}{(n-1)(n-2) \dots (n-r+1)} \right\}_{n=1}^\infty$$

is nonincreasing, and so it is converging, completing the proof of (6) and (8) for $\alpha > 1$. $\square$

Proof of Proposition 10 For every $G \in \mathcal{P}$, Proposition 9 gives

$$\left(\lambda^{(\beta)}(G) \right)^\beta \leq \left(\lambda^{(\alpha)}(G) \right)^\alpha (r!e(G))^{\beta-\alpha}.$$

Hence, choosing $G \in \mathcal{P}_n$ such that $\lambda^{(\beta)}(G) = \lambda^{(\beta)}(\mathcal{P}, n)$, we find that

$$\begin{aligned} \left(\frac{\lambda^{(\beta)}(\mathcal{P}, n) n^{r/\beta-1}}{(n-1)_{r-1}} \right)^\beta &= \left(\frac{\lambda^{(\beta)}(G) n^{r/\beta-1}}{(n-1)_{r-1}} \right)^\beta \\ &\leq \left(\frac{\lambda^{(\beta)}(G) n^{r/\beta-1}}{(n-1)_{r-1}} \right)^\alpha \left(\frac{r!e(G)}{(n)_r} \right)^{\beta-\alpha} \\ &\leq \left(\frac{\lambda^{(\beta)}(G) n^{r/\beta-1}}{(n-1)_{r-1}} \right)^\alpha \left(ex(\mathcal{P}, n) / \binom{n}{r} \right)^{\beta-\alpha}. \end{aligned}$$

Letting now $n \rightarrow \infty$, we see that

$$\left(\lambda^{(\beta)}(\mathcal{P}) \right)^\beta \leq \left(\lambda^{(\alpha)}(\mathcal{P}) \right)^\alpha (\pi(\mathcal{P}))^{\beta-\alpha}$$

completing the proof of (10). $\square$

Proof of Theorem 11 For the purposes of this proof let us write $k_H(G)$ for the number of subgraphs of G which are isomorphic to H .

We start by recalling the Hypergraph Removal Lemma, one of the most important consequences of the Hypergraph Regularity Lemma, proved independently by Gowers [11] and by Nagle, Rödl, Schacht and Skokan [23], [35].

Removal Lemma *Let H be an r -graph of order h and let $\varepsilon > 0$. There exists $\delta = \delta_H(\varepsilon) > 0$ such that if G is an r -graph of order n , with $k_H(G) < \delta n^h$, then there is an r -graph $G_0 \subset G$ such that $e(G_0) \geq e(G) - \varepsilon n^r$ and $k_H(G_0) = 0$.*

In [6] Erdős showed that for every $\varepsilon > 0$ there exists $\delta > 0$ such that if G is an r -graph with $e(G) \geq \varepsilon n^r$, then $K_r(k, \dots, k) \subset G$ for some $k \geq \delta (\log n)^{1/(r-1)}$. As noted by Rödl and Schacht [36] (also by Bollobás, unpublished) this result of Erdős implies the following general assertion.

Theorem A *Let H be an r -graph of order h and let $\varepsilon > 0$. There exists $\delta = \delta_H(\varepsilon) > 0$ such that if G is an r -graph of order n , with $k_H(G) \geq \varepsilon n^h$, then $H(k, \dots, k) \subset G$ for some $k = \left\lceil \delta (\log n)^{1/(h-1)} \right\rceil$.*

Suppose now that H is an r -graph of order h , let $H(k_1, \dots, k_h)$ be a fixed blow-up of H , and set $k = \max\{k_1, \dots, k_h\}$. Take $G \in \text{Mon}(H(k_1, \dots, k_h))_n$ such that

$$\lambda^{(\alpha)}(G) = \lambda^{(\alpha)}(\text{Mon}(H(k_1, \dots, k_h)), n)$$

For every $\varepsilon > 0$, choose $\delta = \delta_H(\varepsilon)$, as in the Removal Lemma. Since $H(k, \dots, k) \not\subset G$, Theorem A implies that if n is sufficiently large, then $k_H(G) < \delta n^h$. Now the Removal Lemma implies that there is an r -graph $G_0 \subset G$ such that $e(G_0) \geq e(G) - \varepsilon n^r$ and $k_H(G_0) = 0$. Clearly, we can assume that $V(G_0) = V(G)$. By Proposition 29, we see that

$$\lambda^{(\alpha)}(G) \leq \lambda^{(\alpha)}(G_0) + (\varepsilon r! n^r)^{1-1/\alpha},$$

and hence,

$$\begin{aligned} \frac{\lambda^{(\alpha)}(\text{Mon}(H(k, \dots, k)), n) n^{r/\alpha-1}}{(n-1)_{r-1}} &\leq \frac{(\lambda^{(\alpha)}(G_0) + (\varepsilon r! n^r)^{1-1/\alpha}) n^{r/\alpha-1}}{(n-1)_{r-1}} \\ &\leq \lambda^{(\alpha)}(\text{Mon}(H)) + o(1) + \frac{(\varepsilon r!)^{1-1/\alpha} n^{r-r/\alpha} n^{1-r/\alpha}}{n^{r-1}} \\ &= \lambda^{(\alpha)}(\text{Mon}(H)) + o(1) + (\varepsilon r!)^{1-1/\alpha}. \end{aligned}$$

Since ε can be made arbitrarily small, we see that

$$\lambda^{(\alpha)}(\text{Mon}H(k_1, \dots, k_h)) \leq \lambda^{(\alpha)}(\text{Mon}(H)),$$

completing the proof of Theorem 11. □

Proof Theorem 9 Let $1 \leq \alpha < \beta$. Using (10), after some cancellations, we find that

$$\lambda^{(\beta)}(\mathcal{P}) \leq \left(\lambda^{(\alpha)}(\mathcal{P}) \right)^{\alpha/\beta} (\pi(\mathcal{P}))^{1-\alpha/\beta}.$$

From (9) we have $\pi(\mathcal{P}) \leq \lambda^{(\alpha)}(\mathcal{P})$ and so

$$(\pi(\mathcal{P}))^{1-\alpha/\beta} \leq \left(\lambda^{(\alpha)}(\mathcal{P}) \right)^{1-\alpha/\beta}.$$

Substituting in the above, we see that

$$\lambda^{(\beta)}(\mathcal{P}) \leq \left(\lambda^{(\alpha)}(\mathcal{P}) \right)^{\alpha/\beta} \left(\lambda^{(\alpha)}(\mathcal{P}) \right)^{1-\alpha/\beta} = \lambda^{(\alpha)}(\mathcal{P}),$$

proving Theorem 9. □

Proof of Theorem 12 As mentioned above, if $\lambda(\mathcal{P}) = \pi(\mathcal{P})$, then

$$\lambda^{(\alpha)}(\mathcal{P}) = \pi(\mathcal{P})$$

for every $\alpha > r$. Thus, all we need to prove is the case $1 < \alpha \leq r$. Fix α in the indicated range, and for every natural n , set for short $\lambda_n = \lambda^{(\alpha)}(\mathcal{P}, n)$. Note that if $\lambda^{(\alpha)}(\mathcal{P}) = 0$, then inequality (9) implies that $\pi(\mathcal{P}) = 0$; therefore Theorem 12 holds for $\lambda^{(\alpha)}(\mathcal{P}) = 0$. We shall assume hereafter that $\lambda^{(\alpha)}(\mathcal{P}) > 0$. Recall also that $(n)_r$ stands for $n!/(n-r)!$.

Claim A *There are infinitely many n for which*

$$\frac{\lambda_{n-1}(n-1)^{r/\alpha-1}}{(n-2)_{r-1}} - \frac{\lambda_n n^{r/\alpha-1}}{(n-1)_{r-1}} < \frac{1}{n \log n} \cdot \frac{\lambda_n n^{r/\alpha-1}}{(n-1)_{r-1}}$$

Proof Indeed, assume for a contradiction that there is n_0 such that

$$\frac{\lambda_{n-1}(n-1)^{r/\alpha-1}}{(n-2)_{r-1}} - \frac{\lambda_n n^{r/\alpha-1}}{(n-1)_{r-1}} \geq \frac{1}{n \log n} \cdot \frac{\lambda_n n^{r/\alpha-1}}{(n-1)_{r-1}}$$

for every $n > n_0$. Then for every $k > n_0$ we see that

$$\begin{aligned} \frac{\lambda_{n_0-1}(n_0-1)^{r/\alpha-1}}{(n_0-1)_{r-1}} - \frac{\lambda_k k^{r/\alpha-1}}{(k-1)_{r-1}} &= \sum_{n=n_0}^k \frac{\lambda_{n-1}(n-1)^{r/\alpha-1}}{(n-2)_{r-1}} - \frac{\lambda_n n^{r/\alpha-1}}{(n-1)_{r-1}} \\ &\geq \sum_{n=n_0}^k \frac{1}{n \log n} \frac{\lambda_n n^{r/\alpha-1}}{(n-1)_{r-1}} \\ &\geq \lambda^{(\alpha)}(\mathcal{P}) \sum_{n=n_0}^k \frac{1}{n \log n}. \end{aligned}$$

This is a contradiction, since the left-hand side is bounded and the right-hand side diverges, proving Claim A.

Using Claim A, choose sufficiently large n so that

$$\frac{\lambda_{n-1}(n-1)^{r/\alpha-1}}{(n-2)_{r-1}} - \frac{\lambda_n n^{r/\alpha-1}}{(n-1)_{r-1}} < \frac{1}{n \log n} \cdot \frac{\lambda_n n^{r/\alpha-1}}{(n-1)_{r-1}}.$$

After some rearrangement we obtain

$$\frac{\lambda_{n-1}}{\lambda_n} < \frac{n^{r/\alpha-1}(n-r)}{(n-1)^{r/\alpha}} \left(1 + \frac{1}{n \log n}\right). \quad (39)$$

Let $G \in \mathcal{P}_n$ be such that $\lambda^{(\alpha)}(G) = \lambda_n$ and let $\mathbf{x} = (x_1, \dots, x_n)$ be a nonnegative vector satisfying $\|\mathbf{x}\|_\alpha = 1$ and

$$\lambda_n = \lambda^{(\alpha)}(G) = P_G(\mathbf{x}).$$

Let $k \in V(G)$ be a vertex and let $\mathbf{x}'$ be the $(n-1)$ -vector obtained from $\mathbf{x}$ by omitting x_k . For the graph $G - k$ we have

$$\begin{aligned} P_{G-k}(\mathbf{x}') &= P_G(\mathbf{x}) - r!x_k \sum_{\{k, i_1, \dots, i_{r-1}\} \in E(G)} x_{i_1} \dots x_{i_{r-1}} \\ &= \lambda^{(\alpha)}(G) - rx_k \left(\lambda^{(\alpha)}(G) x_k^{\alpha-1} \right) = \lambda_n (1 - rx_k^\alpha) \end{aligned} \quad (40)$$

On the other hand, $\mathcal{P}$ is a hereditary property and so $G - k \in \mathcal{P}_{n-1}$. Therefore,

$$P_{G-k}(\mathbf{x}') \leq \lambda(G-k) \|\mathbf{x}'\|_\alpha^r = \lambda(G-k) (1 - x_k^\alpha)^{r/\alpha} \leq \lambda_{n-1} (1 - x_k^\alpha)^{r/\alpha}.$$

This inequality, together with (40), implies that

$$\lambda_n (1 - rx_k^\alpha) \leq \lambda_{n-1} (1 - x_k^\alpha)^{r/\alpha}.$$

Hence, in view of (39),

$$\frac{1 - rx_k^\alpha}{(1 - x_k^\alpha)^{r/\alpha}} \leq \frac{\lambda_{n-1}}{\lambda_n} \leq \frac{n^{r/\alpha-1}(n-r)}{(n-1)^{r/\alpha}} \left(1 + \frac{1}{n \log n}\right). \quad (41)$$

We shall prove that x_k^α is sufficiently large to apply Lemma 32.

Claim B For n sufficiently large,

$$x_k^\alpha > \frac{1}{n} \left(1 - \frac{1}{(\alpha-1) \log n}\right).$$

Proof Indeed, assume for a contradiction that

$$x_k^\alpha \leq \frac{1}{n} \left(1 - \frac{1}{(\alpha-1) \log n}\right). \quad (42)$$

Note that the function

$$f(y) = \frac{1 - ry}{(1 - y)^{r/\alpha}}$$

is decreasing in y for $0 \leq y < 1$, because

$$\begin{aligned} \frac{df(y)}{dy} &= \frac{-r(1-y)^{r/\alpha} + \frac{r}{\alpha}(1-ry)(1-y)^{r/\alpha-1}}{(1-y)^{2r/\alpha}} \\ &= \frac{r(1-y)^{r/\alpha-1}}{\alpha(1-y)^{2r/\alpha}} (-\alpha(1-y) + 1-ry) \\ &= \frac{r(1-y)^{r/\alpha-1}}{\alpha(1-y)^{2r/\alpha}} (-(\alpha-1) - (r-1)y) \leq 0. \end{aligned}$$

Hence, (42) implies that

$$\frac{1 - \frac{r}{n} \left(1 - \frac{1}{(\alpha-1)\log n}\right)}{1 - \frac{1}{n} \left(1 - \frac{1}{(\alpha-1)\log n}\right)^{r/\alpha}} = f\left(\frac{1}{n} \left(1 - \frac{1}{(\alpha-1)\log n}\right)\right) \leq f(x_k^\alpha) = \frac{1 - rx_k^\alpha}{(1 - x_k^\alpha)^{r/\alpha}}.$$

Combining this inequality with (41), we see that

$$\frac{1 - \frac{r}{n} \left(1 - \frac{1}{(\alpha-1)\log n}\right)}{1 - \frac{1}{n} \left(1 - \frac{1}{(\alpha-1)\log n}\right)^{r/\alpha}} \leq \frac{n^{r/\alpha-1}(n-r)}{(n-1)^{r/\alpha}} \left(1 + \frac{1}{n \log n}\right),$$

and so,

$$\frac{\left(n - r + \frac{r}{(\alpha-1)\log n}\right) n^{r/\alpha-1}}{\left(n - 1 + \frac{1}{(\alpha-1)\log n}\right)^{r/\alpha}} = \frac{1 - \frac{r}{n} \left(1 - \frac{1}{(\alpha-1)\log n}\right)}{\left(1 - \frac{1}{n} \left(1 - \frac{1}{(\alpha-1)\log n}\right)\right)^{r/\alpha}} \leq \frac{n^{r/\alpha-1}(n-r)}{(n-1)^{r/\alpha}} \left(1 + \frac{1}{n \log n}\right).$$

Rearranging this inequality, we obtain

$$1 + \frac{r}{(\alpha-1)(n-r)\log n} \leq \left(1 + \frac{1}{(\alpha-1)(n-1)\log n}\right)^{r/\alpha} \left(1 + \frac{1}{n \log n}\right) \quad (43)$$

To simplify the right-hand side of (43), using Bernoulli's inequality, we obtain for n sufficiently large,

$$\begin{aligned} \left(1 + \frac{1}{(\alpha-1)(n-1)\log n}\right)^{r/\alpha} &= \frac{1}{\left(1 - \frac{1}{(\alpha-1)(n-1)\log n+1}\right)^{r/\alpha}} \leq \frac{1}{1 - \frac{r/\alpha}{(\alpha-1)(n-1)\log n+1}} \\ &= 1 + \frac{r/\alpha}{(\alpha-1)(n-1)\log n - r/\alpha + 1} \\ &\leq 1 + \frac{r/\alpha}{(\alpha-1)(n-2)\log n}. \end{aligned}$$

Hence,

$$\begin{aligned} 1 + \frac{r}{(\alpha-1)(n-r)\log n} &\leq \left(1 + \frac{r/\alpha}{(n-2)(\alpha-1)\log n}\right) \left(1 + \frac{1}{n \log n}\right) \\ &\leq 1 + \frac{r/\alpha}{(n-2)(\alpha-1)\log n} + \frac{1}{n \log n} + \frac{r/\alpha}{(\alpha-1)n(n-2)\log^2 n}. \end{aligned}$$

After some cancellations and rearranging, we obtain

$$\frac{r\alpha}{n-r} \leq \frac{r}{n-2} + \frac{(\alpha-1)\alpha}{n} + \frac{r}{n(n-2)\log n}.$$

Now in view of $r \geq 2$ and $\alpha \leq r$, we see that

$$\begin{aligned} \frac{r\alpha}{n-2} &\leq \frac{r\alpha}{n-r} \leq \frac{r}{n-2} + \frac{(\alpha-1)\alpha}{n} + \frac{r}{n(n-2)\log n} \\ &\leq \frac{r}{n-2} + \frac{(\alpha-1)r}{n} + \frac{r}{n(n-2)\log n}, \end{aligned}$$

finally reducing to

$$2(\alpha-1) \leq \frac{1}{\log n},$$

which is a contradiction for n large. This completes the proof of Claim B.

Hence, if n is sufficiently large, for every $k \in V(G)$, we have

$$x_k^r > \frac{1}{n} \left(1 - \frac{1}{(\alpha-1)\log n} \right),$$

and so, Lemma 32 implies that

$$\frac{\lambda_n n^{r/\alpha-1}}{(r-1)!} \leq \delta + \frac{2r}{\alpha(\alpha-1)\log n} \binom{n-1}{r-1} \leq \frac{re(G)}{n} + \frac{2r}{\alpha(\alpha-1)\log n} \binom{n-1}{r-1}.$$

Therefore,

$$\frac{\lambda_n n^{r/\alpha-1}}{(n-1)_{r-1}} \leq \frac{e(G)}{\frac{n}{r} \binom{n-1}{r-1}} + \frac{2r}{\alpha(\alpha-1)\log n} \leq \frac{ex(\mathcal{P}, n)}{\binom{n}{r}} + \frac{2r}{\alpha(\alpha-1)\log n}.$$

Since n can be arbitrary large, we can pass to limits obtaining

$$\lambda^{(\alpha)}(\mathcal{P}) \leq \pi(\mathcal{P}).$$

This inequality together with (13) completes the proof of Theorem 12. $\square$

Proof of Theorem 15 Since Proposition 1 implies that $\lambda^{(1)}(\mathcal{P}) \geq \pi(\mathcal{P})$, to finish the proof we shall show that $\lambda^{(1)}(\mathcal{P}) \leq \pi(\mathcal{P})$. We claim that if $\mathbf{x} = (x_1, \dots, x_n)$ is a nonnegative vector with $\|\mathbf{x}\|_1 = 1$, then

$$P_G(\mathbf{x}) \leq \pi(\mathcal{P}). \quad (44)$$

Because $P_G(\mathbf{x})$ is continuous in each variable, it suffices to prove the inequality for positive rational $x_1, \dots, x_n$. Let thus

$$x_1 = k_1/p, \dots, x_n = k_n/p,$$

where $k_1, \dots, k_n$ are positive integers and p is a common denominator of $x_1, \dots, x_n$. Obviously the condition $x_1 + \dots + x_n = 1$ implies that $p = k_1 + \dots + k_n$. Therefore, (44) it is equivalent to

$$\frac{1}{(k_1 + \dots + k_n)^r} P_G((k_1, \dots, k_n)) \leq \pi(\mathcal{P}). \quad (45)$$

Fix thus positive integers $k_1, \dots, k_n$ and note that

$$P_G((k_1, \dots, k_n)) = P_{G(k_1, \dots, k_n)}((1, \dots, 1)) = r!e(G(k_1, \dots, k_n)).$$

On the other hand, $v(G(k_1, \dots, k_n)) = k_1 + \dots + k_n$ and so

$$e(G(k_1, \dots, k_n)) \leq (\pi(\mathcal{P}) + o(1)) \frac{(k_1 + \dots + k_n)^r}{r!}.$$

Here the term $o(1)$ tends to 0 when $k_1 + \dots + k_n \rightarrow \infty$. Likewise, for every positive integer L , we see that

$$\begin{aligned} \frac{1}{(k_1 + \dots + k_n)^r} P_G((k_1, \dots, k_n)) &= \frac{1}{(Lk_1 + \dots + Lk_n)^r} P_G((Lk_1, \dots, Lk_n)) \\ &= \frac{r!e(G(Lk_1, \dots, Lk_n))}{(Lk_1 + \dots + Lk_n)^r} \leq \pi(\mathcal{P}) + o(1). \end{aligned}$$

Now, letting $L \rightarrow \infty$, we obtain ((45)), and so $\lambda^{(1)}(\mathcal{P}) \leq \pi(\mathcal{P})$, completing the proof of Theorem 15. $\square$

Proof of Theorem 16 Our proof follows an idea of Sidorenko [37], which he used in a similar setting. Let $\mathcal{P}$ be a hereditary and multiplicative family of r -graphs. If $G \in \mathcal{P}_n$, then for every integer $k \geq 1$, we have

$$v(G(k, \dots, k)) = kv(G) \quad \text{and} \quad e(G(k, \dots, k)) = k^r e(G).$$

Therefore,

$$e(G) = \frac{e(G(k, \dots, k))}{k^r} \leq \frac{ex(\mathcal{P}, nk)}{k^r}.$$

Since

$$\lim_{k \rightarrow \infty} \frac{ex(\mathcal{P}, nk)}{\binom{nk}{r}} = \pi(\mathcal{P}),$$

we see that

$$e(G) \leq \pi(\mathcal{P}) \binom{nk}{r} \frac{1}{k^r} + o(k) \binom{nk}{r} \frac{1}{k^r}.$$

Letting $k \rightarrow \infty$, we obtain

$$e(G) \leq \pi(\mathcal{P}) \frac{n^r}{r!},$$

which proves (17).

To prove (18) note that by Proposition 28,

$$\begin{aligned} \lambda^{(\alpha)}(G) &= \lambda^{(\alpha)}(G(k, \dots, k)) k^{r/\alpha - r} \leq \lambda_{nk}^{(\alpha)} k^{r/\alpha - r} \\ &= \left(\lambda^{(\alpha)}(\mathcal{P}) + o(1) \right) (kn)^{r/\alpha - r} \frac{k^{r/\alpha - r}}{(kn)^{r/\alpha - 1}} \\ &= \left(\lambda^{(\alpha)}(\mathcal{P}) + o(1) \right) n^{r - r/\alpha}. \end{aligned}$$

Letting $k \rightarrow \infty$, we see that

$$\lambda^{(\alpha)}(G) \leq \lambda^{(\alpha)}(\mathcal{P}) n^{r - r/\alpha} = \pi(\mathcal{P}) n^{r - r/\alpha},$$

completing the proof of Theorem 16. $\square$

Proof of Theorem 19 If $G \in \mathcal{P}_n$, Proposition 6 implies that

$$\left(\lambda^{(\alpha)}(G)\right)^\alpha \leq \lambda^{(1)}(G) (r!e(G))^{\alpha-1}.$$

Since $\mathcal{P}$ is hereditary and multiplicative, by Theorem 8 we also have

$$\lambda^{(1)}(G) \leq \lambda^{(1)}(\mathcal{P}) = \pi(\mathcal{P}),$$

and so,

$$\lambda^{(1)}(G) \leq \pi(\mathcal{P})^{1/\alpha} (r!e(G))^{1-1/\alpha},$$

completing the proof of Theorem 19. $\square$

Proof of Theorem 13 Let $\mathcal{P}$ be a monotone property of 2-graphs, let $H \notin \mathcal{P}$ be a graph with $\chi(H) = r = \min \{\chi(G) : G \notin \mathcal{P}\}$, and let $K_r(k, \dots, k)$ be the smallest regular r -partite graph containing H . First, obviously $\mathcal{P} \subset \text{Mon}(K_r(k, \dots, k))$, and so, $\lambda^{(\alpha)}(\mathcal{P}) \leq \lambda^{(\alpha)}(\text{Mon}(K_r(k, \dots, k)))$; however, Theorem 11 implies that

$$\lambda^{(\alpha)}(\mathcal{P}) \leq \lambda^{(\alpha)}(\text{Mon}(K_r(k, \dots, k))) = \lambda^{(\alpha)}(\text{Mon}(K_r)) = \frac{r-2}{r-1}.$$

Since $T_{r-1}(n)$, the $(r-1)$ -partite Turán graph of order n , belongs to $\mathcal{P}$, we see that

$$\lambda^{(\alpha)}(\mathcal{P}) = \frac{r-2}{r-1},$$

proving Theorem 13. $\square$

5 Concluding remarks

We have started above a systematic study of the parameter $\lambda^{(\alpha)}(G)$ and its connections to extremal problems for r -graphs. Similarly to eigenvalues of 2-graphs, one may consider other critical points of P_G , for instance, for every r -graph G of order n and every real number $\alpha \geq 1$, define

$$\lambda_{\min}^{(\alpha)}(G) = \min_{\|\mathbf{x}\|_\alpha=1} P_G(\mathbf{x}) = \min_{|x_1|^\alpha + \dots + |x_n|^\alpha = 1} r! \sum_{\{i_1, i_2, \dots, i_r\} \in E(G)} x_{i_1} x_{i_2} \dots x_{i_r}.$$

Obviously $\lambda_{\min}^{(\alpha)}(G)$ is analogous to the smallest eigenvalue of 2-graphs and one can come up with a lot of supporting material about it, including a system of equations similar to (25). In particular, if $\mathcal{P}$ is a hereditary property of r -graphs, we can define

$$\lambda_{\min}^{(\alpha)}(\mathcal{P}, n) = \min_{G \in \mathcal{P}_n} \lambda_{\min}^{(\alpha)}(G),$$

and prove the following statement:

Theorem 33 *Let $\alpha > 1$. If $\mathcal{P}$ is a hereditary property of r -graphs, then the limit*

$$\lambda_{\min}^{(\alpha)}(\mathcal{P}) = \lim_{n \rightarrow \infty} \lambda_{\min}^{(\alpha)}(\mathcal{P}, n) n^{r/\alpha - r}$$

exists.

Many obvious problems arise here, of which we mention the following two:

Problem 1 Let $\alpha > 1$. For a 2-graph G of order n , study its “ α -eigenvalues”, that is to say, critical values of P_G over the unit sphere $|x_1|^\alpha + \cdots + |x_n|^\alpha = 1$.

Since the smallest eigenvalue of 2-graphs has proved to be a useful structural parameter (see, e.g. [25]), one can investigate what role plays $\lambda_{\min}^{(\alpha)}(G)$ for r -graphs.

Problem 2 Let $\alpha > 1$. For an r -graph G of order n , study which structural properties of G are related to $\lambda_{\min}^{(\alpha)}(G)$.

Although for $\alpha \neq 1$ and $\alpha \neq r$ the parameter $\lambda^{(\alpha)}(G)$ is mostly auxiliary, it is challenging and instructive to extend known results about $\lambda^{(r)}(G)$ to general $\alpha \geq 1$. This is interesting even for $r = 2$. Keevash, Lenz and Mubayi have pointed to such generalization in their Corollary 2, but a lot more work is pending. An important initial endeavour would be to recover parts of the Perron-Frobenius theory for $\lambda^{(\alpha)}(G)$. In particular, what is the set of all critical vectors corresponding to $\lambda^{(\alpha)}(G)$.

Problem 3 Given an r -graph G of order n , determine the set of all n -vectors x with $\|x\|_\alpha = 1$ and $\lambda^{(\alpha)}(G) = P_G(x)$.

In view of the importance of flat properties, the following problem is natural, although probably quite difficult:

Problem 4 Characterize all flat properties of r -graphs.

A particular case of this problem arises in connection to Theorem 23. It is curious how rich can be properties of graphs defined by forbidden induced blowups of covering graphs.

Problem 5 For which non-covering graphs F , the property $\text{Her}(\{F\})$ can be represented as $\text{Her}(X)$ for some family X of blowups of covering graphs:?

Acknowledgement Much of the material presented above was conceived in the Spring of 2012, when the author read a course on spectra of hypergraphs at the Math. Department of the University of Memphis. The author is grateful to Peter Keevash, John Lenz and Dhruv Mubayi for sharing their manuscript [16], which greatly motivated the writing of the present paper.

Also, the author is much indebted to Alex Sidorenko for useful discussions. His comments on an earlier version of the paper led to considerable improvement of the presentation.

References

- [1] W. G. Brown, M. Simonovits, Digraph extremal problems, hypergraph extremal problems, and the densities of graph structures, *Disc. Math.* **48** (1984), 147-162
- [2] K. C. Chang, K. Pearson and T. Zhang, Perron-Frobenius theorem for nonnegative hypermatrices, *Commun. Math. Sci.* **6** (2008), 507-520.
- [3] K.C. Chang, K. Pearson and T. Zhang, On eigenvalue problems of real symmetric hypermatrices, *J. Math. Anal. Appl.* **350** (2009) 416-422.
- [4] J. Cooper and A Dutle, Spectra of uniform hypergraphs, *Linear Algebra Appl.* **436** (2012), 3268-3292.

- [5] D. Cvetković, Chromatic number and the spectrum of a graph, *Publ. Inst. Math. (Beograd)* **14(28)** (1972), 25-38, .
- [6] P. Erdős, Extremal problems of graphs and generalized graphs, *Israel J. Math.* **2** (1964), 183-190.
- [7] P. Frankl and V. Rödl, Hypergraphs do not jump, *Combinatorica* **4** (1984), 149-159.
- [8] P. Frankl and Z. Füredi, Extremal problems and the Lagrange function of hypergraphs, *Bulletin Institute Math. Academia Sinica* **16** (1988), 305-313.
- [9] S. Friedland, S. Gaubert and L. Han, Perron-Frobenius theorem for nonnegative multilinear forms, *Linear Algebra Appl.* **438** (2013), 738-749.
- [10] Z. Füredi and M. Simonovits, Triple systems not containing a Fano configuration, *Combin. Probab. Comput.* **14** (2005), 467-484.
- [11] W.T. Gowers, Hypergraph regularity and the multidimensional Szemerédi theorem, *Ann. of Math.* **166** (2007), 897-946.
- [12] G.H. Hardy, J.E. Littlewood, and G. Pólya, *Inequalities, 2nd edition*, Cambridge University Press, 1988, vi+324 pp.
- [13] G. Katona, T. Nemetz and M. Simonovits, On a problem of Turán in the theory of graphs, *Mat. Lapok* **15** (1964), 228-238.
- [14] P. Keevash and B. Sudakov, The Turán number of the Fano plane, *Combinatorica* **25** (2005), 561-574.
- [15] P. Keevash, Hypergraph Turán problems, *Surveys in Combinatorics*, Cambridge University Press, 2011, 83-140.
- [16] P. Keevash, J. Lenz, and D. Mubayi, Spectral extremal problems for hypergraphs, *preprint available at arXiv:1304.0050*.
- [17] N. Khadzhiivanov, Inequalities for graphs (in Russian), *C. R. Acad. Sci. Bul.* **30** (1977), 793-796.
- [18] M.A. Krasnoselskii, Stability of critical values of even functionals on the sphere (in Russian), *Matem. Sb.*, **37:2** (1955), 301-322.
- [19] L.H. Lim, Singular values and eigenvalues of hypermatrices: a variational approach, in *Proceedings of the IEEE International Workshop on Computational Advances in Multi-Sensor Adaptive Processing (CAMSAP '05)* **1** (2005), pp. 129-132.
- [20] L. Lusternik and L. Schnirelman, Topological methods in variational problems (in Russian), *Inst. Mat. Mech., Moscow State Univ.*, 1930.
- [21] L. Lusternik and L. Schnirelman, Méthodes topologiques dans les problèmes variationnels, Paris, Hermann, 1934.
- [22] T. Motzkin and E. Straus, Maxima for graphs and a new proof of a theorem of Turán, *Canad. J. Math.*, **17** (1965), 533-540.
- [23] B. Nagle, V. Rodl and M. Schacht, The counting lemma for regular k-uniform hypergraphs, *Random Structures Algorithms* **28** (2006), 113-179.

- [24] V. Nikiforov, Some inequalities for the largest eigenvalue of a graph. *Combin. Probab. Comput.* **11** (2002), 179–189.
- [25] V. Nikiforov, Bounds on graph eigenvalues II, *Linear Algebra Appl.* **427** (2007), 183–189.
- [26] V. Nikiforov, A spectral condition for odd cycles, *Linear Algebra Appl.* **428** (2008), 1492–1498.
- [27] V. Nikiforov, Some new results in extremal graph theory, *Surveys in Combinatorics*, Cambridge University Press, 2011, 141–181.
- [28] V. Nikiforov, The number of cliques in graphs of given order and size. *Transactions AMS*, **363** (2011), 1599–1618.
- [29] V. Nikiforov, An extension of Maclauren’s inequalities, preprint available at *ArXiv math/0608199*.
- [30] V. Nikiforov, The largest eigenvalue of uniform hypergraphs: basic properties, in preparation.
- [31] V. Nikiforov, Some extremal problems for hereditary properties of graphs, preprint available at *arXiv:1305.1072v1*.
- [32] L. Qi, Eigenvalues of a real supersymmetric hypermatrix, *J. Symbolic Comput.* **40** (2005) 1302–1324.
- [33] L. Qi, Rank and eigenvalues of a supersymmetric hypermatrix, a multivariate homogeneous polynomial and an algebraic surface defined by them, *J. Symbolic Comput.* **41** (2006) 1309–1327.
- [34] L. Qi, Eigenvalues and invariants of hypermatrices, *J. Math. Anal. Appl.* **325** (2007) 1363–1377.
- [35] V. Rodl and J. Skokan, Regularity lemma for uniform hypergraphs, *Random Structures Algorithms* **25** (2004), 1–42.
- [36] V. Rodl and M. Schacht, Complete partite subgraphs in dense hypergraphs, *Random Structures Algorithms*, **41** (2012), 557–573.
- [37] A.F. Sidorenko, On the maximal number of edges in a uniform hypergraph with no forbidden subgraphs (in Russian), *Mat. Zametki*, **41** (1987), 433–455; (English translation in *Math Notes* **41** (1987), 247–259.)
- [38] V. Sós, E. Straus, Extremals of functions on graphs with applications to graphs and hypergraphs, *J. Combin. Theory Ser. B* **32** (1982), 246–257.
- [39] H.S. Wilf, Spectral bounds for the clique and independence numbers of graphs, *J. Combin. Theory Ser. B* **40** (1986), 113–117.
- [40] Y. Yang and Q. Yang, Further results for Perron-Frobenius Theorem for nonnegative tensors, *SIAM Journal on Matrix Analysis*, **31** (2010), 2517–2530.
- [41] Q. Yang and Y. Yang, Further results for Perron-Frobenius Theorem for nonnegative tensors II, *SIAM Journal on Matrix Analysis*, **32** (2011), 1236–1250.
- [42] Y. Yang and Q. Yang, Geometric simplicity of the spectral radius of nonnegative irreducible tensors, *Frontiers of Mathematics in China* **8** (2013), 129–140.